

**ELSEVIER**Contents lists available at ScienceDirect

Journal of Functional Analysis

www.elsevier.com/locate/jfa

The spatial Rokhlin property for actions of compact quantum groups

Selçuk Barlak^a, Gábor Szabó^{b,*}, Christian Voigt^c^a Department of Mathematics and Computer Science, University of Southern Denmark, Campusvej 55, DK-5230 Odense M, Denmark^b Fraser Noble Building, Institute of Mathematics, University of Aberdeen, Aberdeen AB24 3UE, United Kingdom^c School of Mathematics and Statistics, University of Glasgow, 15 University Gardens, Glasgow G12 8QW, United Kingdom

ARTICLE INFO

Article history:

Received 2 June 2016

Accepted 29 September 2016

Available online 8 October 2016

Communicated by Stefaan Vaes

MSC:

primary 46L55, 20G42

secondary 46L35

Keywords:

Rokhlin property

Quantum groups

Classification of C^* -algebras

Classification of dynamical systems

ABSTRACT

We introduce the spatial Rokhlin property for actions of coexact compact quantum groups on C^* -algebras, generalizing the Rokhlin property for both actions of classical compact groups and finite quantum groups. Two key ingredients in our approach are the concept of sequentially split $*$ -homomorphisms, and the use of braided tensor products instead of ordinary tensor products.

We show that various structure results carry over from the classical theory to this more general setting. In particular, we show that a number of C^* -algebraic properties relevant to the classification program pass from the underlying C^* -algebra of a Rokhlin action to both the crossed product and the fixed point algebra. Towards establishing a classification theory, we show that Rokhlin actions exhibit a rigidity property with respect to approximate unitary equivalence. Regarding duality theory, we introduce the notion of spatial approximate representability for actions of discrete quantum groups. The spatial Rokhlin property for actions of a coexact compact quantum group is shown to be dual to spatial approximate

* Corresponding author.

E-mail addresses: barlak@imada.sdu.dk (S. Barlak), gabor.szabo@abdn.ac.uk (G. Szabó), christian.voigt@glasgow.ac.uk (C. Voigt).

representability for actions of its dual discrete quantum group, and vice versa.

© 2016 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Contents

0.	Introduction	2309
1.	Preliminaries	2312
1.1.	Quantum groups	2312
1.2.	Actions, crossed products and Takesaki–Takai duality	2318
1.3.	Braided tensor products	2322
2.	Induced actions on sequence algebras	2323
2.1.	Sequence algebras	2323
2.2.	Induced actions – discrete case	2325
2.3.	Induced actions – compact case	2326
3.	Equivariantly sequentially split $*$ -homomorphisms	2332
4.	The Rokhlin property and approximate representability	2336
4.1.	The spatial Rokhlin property	2336
4.2.	Spatial approximate representability	2340
4.3.	Duality	2343
5.	Rigidity of Rokhlin actions	2346
6.	Examples	2355
	References	2358

0. Introduction

The Rokhlin property for finite group actions on unital C^* -algebras was introduced and studied by Izumi in [15,16], building on earlier work of Herman–Jones [12] and Herman–Ocneanu [13]. Since the very beginning it has proven to be a useful tool in the theory of finite group actions. The Rokhlin property was subsequently generalized by Hirshberg–Winter [14] to the case of compact groups, and studied further by Gardella [9]; see also [11,10]. For finite quantum group actions, Kodaka–Teruya introduced and studied the Rokhlin property and approximate representability in [21].

The established theory, which we shall now briefly summarize in the initial setting of finite group actions, has three particularly remarkable features: The first is a multitude of permanence properties; it is known that many C^* -algebraic properties pass from the coefficient C^* -algebra to the crossed product and fixed point algebra. This was in part addressed by Izumi in [15], and studied more in depth by Osaka–Phillips [27] and Santiago [30]. The second feature is rigidity with respect to approximate unitary equivalence; a result of Izumi [15] asserts that two Rokhlin actions of a finite group on a separable, unital C^* -algebra are conjugate via an approximately inner automorphism if and only if the two actions are pointwise approximately unitarily equivalent; see [11] for the non-unital case and [10] for the case of compact groups. As demonstrated by Izumi in [15], this rigidity property is useful for classifying Rokhlin actions on classifiable C^* -algebras

via K -theoretic invariants. The third feature is duality theory; a result of Izumi [15] shows that an action of a finite abelian group on a separable, unital C^* -algebra has the Rokhlin property if and only if the dual action is approximately representable, and vice versa. This has been generalized to the non-unital case by Nawata [25], and to actions of compact abelian groups by the first two authors [4]; see also [8].

In the present paper, we introduce and study the spatial Rokhlin property for actions of coexact compact quantum groups, generalizing and unifying the work mentioned above. In particular, we carry over various structure results from the classical to the general case. Firstly, this allows us to remove all commutativity assumptions in the study of duality properties for Rokhlin actions. This is relevant even for classical group actions. Indeed, the Pontrjagin dual of a nonabelian group is no longer a group, but can be viewed as a quantum group. Accordingly, a natural way to fully incorporate nonabelian groups into the picture is to work in the setting of quantum groups from the very beginning. Secondly, it turns out that some results can be given quite short and transparent proofs in this more abstract setup, simpler than in previous accounts. Finally, our results are also of interest from the point of view of quantum group theory. Indeed, they provide examples of quantum group actions that either allow for classification, or the systematical analysis of structural properties of crossed product C^* -algebras, in particular whether they fall within the scope of the Elliott program.

Let us highlight two comparably new ingredients in our approach. The first is the notion of (equivariantly) sequentially split $*$ -homomorphisms introduced by the first two authors in [4]. It has already been demonstrated in [4] that many structural results related to the Rokhlin property can be recast and conceptually proved in the language of sequentially split $*$ -homomorphisms, and some new ones could be proved as well. The second ingredient is a purely quantum feature, namely the braided tensor product construction. This provides the correct substitute for tensor product actions in the classical theory, and it also gives a conceptual explanation of the fact that the central sequence algebra is no longer the right tool in the quantum setting. Being widely known in the algebraic theory of quantum groups, braided tensor products in the operator algebraic framework were first introduced and studied in [26].

As already indicated above, for most of the paper we will assume that our quantum groups satisfy exactness/coexactness assumptions. This may appear surprising at first sight; it is essentially due to the fact that we have chosen to work in a reduced setting, that is, with reduced crossed products and minimal (braided) tensor products. We shall indicate at several points in the main text where precisely exactness enters. Our setup yields the strongest versions of conceivable definitions of Rokhlin actions and approximately representable actions, however, at the same time our examples are restricted to the amenable/coamenable case. On the other hand, the reduced setting matches best with the existing literature on quantum group actions, see for instance [2,3,31,26] and references therein. The necessary modifications to set up a full version of our theory are mainly of technical nature; we have refrained from carrying this out here.

Let us now explain how the paper is organized. In Section 1, we gather some preliminaries and background on quantum groups, including a review of Takesaki–Takai duality and braided tensor products. Section 2 deals with induced actions of discrete and compact quantum groups on sequence algebras. Already for classical compact groups these actions typically fail to be continuous, and in the quantum setting this leads to a number of subtle issues. In Section 3, we define and study equivariantly sequentially split $*$ -homomorphisms. We show that, as in the case of group actions, this notion behaves well with respect to crossed products and fixed point algebras. We also establish a general duality result for equivariantly sequentially split $*$ -homomorphisms. In Section 4, we introduce the spatial Rokhlin property for actions of coexact compact quantum groups, and spatial approximate representability for actions of exact discrete quantum groups. We verify that various C^* -algebraic properties pass to crossed products and fixed point algebras. Moreover, we show that the spatial Rokhlin property and spatial approximate representability are dual to each other. In Section 5, we present some steps towards a classification theory for actions with the spatial Rokhlin property. Among other things, we prove that two such G -actions on a C^* -algebra A are conjugate via an approximately inner automorphism if and only if the actions are approximately unitarily equivalent as $*$ -homomorphisms from A to $C^*(G) \otimes A$. This generalizes a number of previous such classification results, in particular those of Izumi [15], Gardella–Santiago [10] and Kodaka–Teruya [21]. In this section, we also generalize a K -theory formula for the fixed-point algebra of a Rokhlin action, first proved for certain finite group cases by Izumi [15] and for compact group actions by the first two authors in [4]. Finally, in Section 6 we present some examples of Rokhlin actions. In particular, we show that any coamenable compact quantum group admits an essentially unique action with the spatial Rokhlin property on \mathcal{O}_2 .

The work presented here was initiated while the first two authors participated in the conference CSTAR at the University of Glasgow in September 2014. Substantial parts of this work were carried out during research visits of the authors to Oberwolfach in June 2015, of the first two authors at the Mittag-Leffler Institute from January to March 2016, and of the third author to the University of Southern Denmark in April 2016. The authors are grateful to all these institutions for their hospitality and support.

The first author was supported by GIF Grant 1137-30.6/2011, ERC AdG 267079, SFB 878 ‘Groups, Geometry, and Actions’ and the Villum Fonden project grant ‘Local and global structures of groups and their algebras’ (2014–2018), 7423. The second author was supported by SFB 878 ‘Groups, Geometry, and Actions’ and the Engineering and Physical Sciences Research Council Grant EP/N00874X/1 ‘Regularity and dimension for C^* -algebras’. The third author was supported by the Engineering and Physical Sciences Research Council Grant EP/L013916/1 and the Polish National Science Centre grant no. 2012/06/M/ST1/00169.

The authors would like to thank the referee for a thorough reading of the manuscript and for helpful suggestions.

1. Preliminaries

In this preliminary section we collect some definitions and results from the theory of quantum groups and fix our notation. We will mainly follow the conventions in [26] as far as general quantum group theory is concerned. For more detailed information and background we refer to [34,24,22,23].

Let us make some general remarks on the notation used throughout the paper. We write $\mathbb{L}(\mathcal{E})$ for the space of adjointable operators on a Hilbert A -module, and $\mathbb{K}(\mathcal{E})$ denotes the space of compact operators. The closed linear span of a subset X of a Banach space is denoted by $[X]$. If x, y are elements of a Banach space and $\varepsilon > 0$ we write $x =_\varepsilon y$ if $\|x - y\| < \varepsilon$.

Depending on the context, the symbol \otimes denotes either the tensor product of Hilbert spaces, the minimal tensor product of C^* -algebras, or the tensor product of von Neumann algebras. We write \odot for algebraic tensor products.

If A and B are C^* -algebras then the flip map $A \otimes B \rightarrow B \otimes A$ is denoted by σ . That is, we have $\sigma(a \otimes b) = b \otimes a$ for $a \in A, b \in B$.

If \mathcal{H} is a Hilbert space we write $\Sigma \in \mathbb{L}(\mathcal{H} \otimes \mathcal{H})$ for the flip map $\Sigma(\xi \otimes \eta) = \eta \otimes \xi$. For operators on multiple tensor products we use the leg numbering notation. For instance, if $W \in \mathbb{L}(\mathcal{H} \otimes \mathcal{H})$ is an operator on $\mathcal{H} \otimes \mathcal{H}$, then $W_{12} = W \otimes \text{id} \in \mathbb{L}(\mathcal{H} \otimes \mathcal{H} \otimes \mathcal{H})$ and $W_{23} = \text{id} \otimes W$. Moreover, $W_{13} = \tilde{\Sigma}_{12} W_{23} \Sigma_{12}$.

If B is a C^* -algebra we write \tilde{B} for the smallest unitarization of B .

1.1. Quantum groups

Although we will only be interested in compact and discrete quantum groups, let us first recall a few definitions and facts regarding general locally compact quantum groups.

Let φ be a normal, semifinite and faithful weight on a von Neumann algebra M . We use the standard notation

$$\mathcal{M}_\varphi^+ = \{x \in M_+ \mid \varphi(x) < \infty\}, \quad \mathcal{N}_\varphi = \{x \in M \mid \varphi(x^*x) < \infty\}$$

and write M_*^+ for the space of positive normal linear functionals on M . Assume that $\Delta : M \rightarrow M \otimes M$ is a normal unital $*$ -homomorphism. The weight φ is called left invariant with respect to Δ if

$$\varphi((\omega \otimes \text{id})\Delta(x)) = \varphi(x)\omega(\mathbf{1})$$

for all $x \in \mathcal{M}_\varphi^+$ and $\omega \in M_*^+$. Similarly one defines the notion of a right invariant weight.

Definition 1.1. A locally compact quantum group G is given by a von Neumann algebra $L^\infty(G)$ together with a normal unital $*$ -homomorphism $\Delta : L^\infty(G) \rightarrow L^\infty(G) \otimes L^\infty(G)$ satisfying the coassociativity relation

$$(\Delta \otimes \text{id})\Delta = (\text{id} \otimes \Delta)\Delta$$

and normal semifinite faithful weights φ and ψ on $L^\infty(G)$ which are left and right invariant, respectively. The weights φ and ψ will also be referred to as Haar weights of G .

Remark 1.2. Our notation should help to make clear how ordinary locally compact groups can be viewed as quantum groups. Indeed, if G is a locally compact group, then the algebra $L^\infty(G)$ of essentially bounded measurable functions on G together with the comultiplication $\Delta : L^\infty(G) \rightarrow L^\infty(G) \otimes L^\infty(G)$ given by

$$\Delta(f)(s, t) = f(st)$$

defines a locally compact quantum group. The weights φ and ψ are given in this case by left and right Haar measures, respectively. Of course, for a general locally compact quantum group G the notation $L^\infty(G)$ is purely formal. Similar remarks apply to the C^* -algebras $C_r^*(G)$, $C_r^*(G)$ and $C_0^f(G)$, $C_0^r(G)$ associated to G that we discuss below. It is convenient to view all of them as different appearances of the quantum group G .

Remark 1.3 (cf. [22]). Let G be a locally compact quantum group and let $\Lambda : \mathcal{N}_\varphi \rightarrow L^2(G)$ be the GNS-construction for the Haar weight φ . Throughout the paper we will only consider second countable quantum groups, that is, quantum groups for which $L^2(G)$ is a separable Hilbert space.

One obtains a unitary $W_G = W$ on $L^2(G) \otimes L^2(G)$ by

$$W^*(\Lambda(x) \otimes \Lambda(y)) = (\Lambda \otimes \Lambda)(\Delta(y)(x \otimes \mathbf{1}))$$

for all $x, y \in \mathcal{N}_\varphi$. This unitary is multiplicative, which means that W satisfies the pentagonal equation

$$W_{12}W_{13}W_{23} = W_{23}W_{12}.$$

From W one can recover the von Neumann algebra $L^\infty(G)$ as the strong closure of the algebra $(\text{id} \otimes \mathbb{L}(L^2(G))_*)(W)$, where $\mathbb{L}(L^2(G))_*$ denotes the space of normal linear functionals on $\mathbb{L}(L^2(G))$. Moreover one has

$$\Delta(x) = W^*(\mathbf{1} \otimes x)W$$

for all $x \in M$. The algebra $L^\infty(G)$ has an antipode, which is an unbounded, σ -strong* closed linear map S given by $S(\text{id} \otimes \omega)(W) = (\text{id} \otimes \omega)(W^*)$ for $\omega \in \mathbb{L}(L^2(G))_*$. Moreover, there is a polar decomposition $S = R\tau_{-i/2}$ where R is an antiautomorphism of $L^\infty(G)$ called the unitary antipode and (τ_t) is a strongly continuous one-parameter group of

automorphisms of $L^\infty(G)$ called the scaling group. The unitary antipode satisfies $\sigma \circ (R \otimes R) \circ \Delta = \Delta \circ R$.

The group-von Neumann algebra $\mathcal{L}(G)$ of the quantum group G is the strong closure of the algebra $(\mathbb{L}(L^2(G))_* \otimes \text{id})(W)$ with the comultiplication $\hat{\Delta} : \mathcal{L}(G) \rightarrow \mathcal{L}(G) \otimes \mathcal{L}(G)$ given by

$$\hat{\Delta}(y) = \hat{W}^*(1 \otimes y)\hat{W}$$

where $\hat{W} = \Sigma W^* \Sigma$ and $\Sigma \in \mathbb{L}(L^2(G) \otimes L^2(G))$ is the flip map. It defines a locally compact quantum group \hat{G} which is called the dual of G . The left invariant weight $\hat{\varphi}$ for the dual quantum group has a GNS-construction $\hat{\Lambda} : \mathcal{N}_{\hat{\varphi}} \rightarrow L^2(G)$, and according to our conventions we have $\mathcal{L}(G) = L^\infty(\hat{G})$.

Remark 1.4. Since we are following the conventions of Kustermans and Vaes [22], there is a flip map built into the definition of $\hat{\Delta}$. As we will see below, this is a natural choice when working with Yetter–Drinfeld actions; however, it is slightly inconvenient when it comes to Takesaki–Takai duality. We will write \check{G} for the quantum group corresponding to $\mathcal{L}(G)^{\text{cop}}$. That is, $L^\infty(\check{G})$ is the von Neumann algebra $\mathcal{L}(G)$ equipped with the opposite comultiplication $\check{\Delta} = \hat{\Delta}^{\text{cop}} = \sigma \circ \hat{\Delta}$, where σ denotes the flip map. By slight abuse of terminology, we shall refer to both \check{G} and \hat{G} as the dual of G . According to Pontrjagin duality, the double dual of G in either of these conventions is canonically isomorphic to G .

Remark 1.5 (cf. [22]). The modular conjugations of the left Haar weights φ and $\hat{\varphi}$ are denoted by J and \hat{J} , respectively. These operators implement the unitary antipodes in the sense that

$$R(x) = \hat{J}x^*\hat{J}, \quad \hat{R}(y) = Jy^*J$$

for $x \in L^\infty(G)$ and $y \in \mathcal{L}(G)$. Note that $L^\infty(G)' = JL^\infty(G)J$ and $\mathcal{L}(G)' = \hat{J}\mathcal{L}(G)\hat{J}$ for the commutants of $L^\infty(G)$ and $\mathcal{L}(G)$. Using J and \hat{J} one obtains multiplicative unitaries

$$V = (\hat{J} \otimes \hat{J})\hat{W}(\hat{J} \otimes \hat{J}), \quad \hat{V} = (J \otimes J)W(J \otimes J)$$

which satisfy $V \in \mathcal{L}(G)' \otimes L^\infty(G)$ and $\hat{V} \in L^\infty(G)' \otimes \mathcal{L}(G)$, respectively. We have

$$\Delta(f) = V(f \otimes 1)V^*, \quad \hat{\Delta}(x) = \hat{V}(x \otimes 1)\hat{V}^*$$

for $f \in L^\infty(G)$ and $x \in \mathcal{L}(G)$. We also record the formula

$$(\hat{J} \otimes J)W(\hat{J} \otimes J) = W^*,$$

which is equivalent to saying $(R \otimes \hat{R})(W) = W$.

We will mainly work with the C^* -algebras associated to the locally compact quantum group G . The algebra $[(\text{id} \otimes \mathbb{L}(L^2(G))_*)(W)]$ is a strongly dense C^* -subalgebra of $L^\infty(G)$ which we denote by $C_0^r(G)$. Dually, the algebra $[(\mathbb{L}(L^2(G))_* \otimes \text{id})(W)]$ is a strongly dense C^* -subalgebra of $\mathcal{L}(G)$ which we denote by $C_r^*(G)$. These algebras are the reduced algebra of continuous functions vanishing at infinity on G and the reduced group C^* -algebra of G , respectively. One has $W \in M(C_0^r(G) \otimes C_r^*(G))$.

Restriction of the comultiplications on $L^\infty(G)$ and $\mathcal{L}(G)$ turns $C_0^r(G)$ and $C_r^*(G)$ into Hopf C^* -algebras in the following sense.

Definition 1.6. A Hopf C^* -algebra is a C^* -algebra H together with an injective nondegenerate $*$ -homomorphism $\Delta : H \rightarrow M(H \otimes H)$ such that the diagram

$$\begin{array}{ccc} H & \xrightarrow{\Delta} & M(H \otimes H) \\ \downarrow \Delta & & \downarrow \text{id} \otimes \Delta \\ M(H \otimes H) & \xrightarrow{\Delta \otimes \text{id}} & M(H \otimes H \otimes H) \end{array}$$

is commutative and $[\Delta(H)(1 \otimes H)] = H \otimes H = [(H \otimes 1)\Delta(H)]$.

Definition 1.7. A compact quantum group is a locally compact quantum group G such that $C_0^r(G)$ is unital. Similarly, a discrete quantum group is a locally compact quantum group G such that $C_r^*(G)$ is unital.

We will write $C^r(G)$ instead of $C_0^r(G)$ if G is a compact quantum group. A finite quantum group is a compact quantum group G such that $C^r(G)$ is finite dimensional. This is the case if and only if $C_r^*(G)$ is finite dimensional.

Definition 1.8. A unitary corepresentation of a Hopf C^* -algebra H on a Hilbert B -module \mathcal{E} is a unitary $U \in M(H \otimes \mathbb{K}(\mathcal{E})) = \mathbb{L}(H \otimes \mathcal{E})$ such that

$$(\Delta \otimes \text{id})(U) = U_{13}U_{23}.$$

A unitary representation of a locally compact quantum group G is a unitary corepresentation of $C_0^r(G)$.

Remark 1.9. This terminology is compatible with the classical case, more precisely, for a classical locally compact group G there is a bijective correspondence between unitary corepresentations of $C_0(G)$ on Hilbert spaces and strongly continuous unitary representations of G in the usual sense, see [24, Section 5] and [31, 5.2.5].

Example 1.10. The (left) regular representation of a locally compact quantum group G is the representation of G on the Hilbert space $L^2(G)$ given by the multiplicative unitary

$W \in M(C_0^r(G) \otimes \mathbb{K}(L^2(G)))$, see [Remark 1.3](#). In fact, the relation $(\Delta \otimes \text{id})(W) = W_{13}W_{23}$ is equivalent to the pentagon equation for W .

Remark 1.11. Let G be a locally compact quantum group. The full group C^* -algebra of G is a Hopf C^* -algebra $C_f^*(G)$ together with a unitary representation $\mathcal{W} \in M(C_0^r(G) \otimes C_f^*(G))$ satisfying the following universal property: for every unitary representation $U \in M(C_0^r(G) \otimes \mathbb{K}(\mathcal{E}))$ of G there exists a unique nondegenerate $*$ -homomorphism $\pi : C_f^*(G) \rightarrow \mathbb{K}(\mathcal{E})$ such that $(\text{id} \otimes \pi)(\mathcal{W}) = U$.

Similarly, one obtains the full C^* -algebra $C_0^f(G)$ of functions on G .

Definition 1.12. Let G be a locally compact quantum group. We say that G is amenable if the canonical quotient map $C_f^*(G) \rightarrow C_r^*(G)$ is an isomorphism. Similarly, G is called coamenable if the dual \hat{G} is amenable.

If G is coamenable we will simply write $C_0(G)$ for $C_0^r(G)$. By slight abuse of notation, we will also write $C_0(G)$ if a statement holds for both $C_0^f(G)$ and $C_0^r(G)$. In particular, if G is compact and coamenable we will simply write $C(G)$ instead $C^r(G)$. Similarly, if G is amenable we will write $C^*(G)$ for $C_r^*(G)$. We remark that every compact quantum group is amenable, and equivalently every discrete quantum group is coamenable.

Remark 1.13 (cf. [\[34, 24\]](#)). Let G be a compact quantum group. In analogy with the theory for compact groups, every unitary representation of G is completely reducible, and all irreducible representations are finite dimensional. We write $\text{lrr}(G)$ for the set of equivalence classes of irreducible unitary representations of G . Our general second countability assumption amounts to saying that the set $\text{lrr}(G)$ is countable.

A unitary representation of G on a finite dimensional Hilbert space \mathcal{H}_λ is given by a unitary $u^\lambda \in C^r(G) \otimes \mathbb{K}(\mathcal{H}_\lambda)$, so it can be viewed as an element $u^\lambda = (u_{ij}^\lambda) \in M_n(C^r(G))$ if $\dim(\mathcal{H}_\lambda) = n$. Moreover, the corepresentation identity translates into the formula

$$\Delta(u_{ij}^\lambda) = \sum_{k=1}^n u_{ik}^\lambda \otimes u_{kj}^\lambda$$

for the comultiplication of the matrix coefficients u_{ij}^λ .

By Peter–Weyl theory, the linear span of all matrix coefficients u_{ij}^λ for $\lambda \in \text{lrr}(G)$ forms a dense $*$ -subalgebra $\mathcal{O}(G) \subset C^r(G)$. In fact, together with the counit $\epsilon : \mathcal{O}(G) \rightarrow \mathbb{C}$ given by

$$\epsilon(u_{ij}^\lambda) = \delta_{ij}$$

and the antipode $S : \mathcal{O}(G) \rightarrow \mathcal{O}(G)$ given by

$$S(u_{ij}^\lambda) = (u_{ji}^\lambda)^*$$

the algebra $\mathcal{O}(G)$ becomes a Hopf $*$ -algebra.

We will use the Sweedler notation $\Delta(f) = f_{(1)} \otimes f_{(2)}$ for the comultiplication of general elements of $\mathcal{O}(G)$. This is useful for bookkeeping of coproducts, let us emphasize however that this notation is not meant to say that $\Delta(f)$ is a simple tensor. For higher coproducts one introduces further indices, for instance $f_{(1)} \otimes f_{(2)} \otimes f_{(3)}$ is an abbreviation for $(\Delta \otimes \text{id})\Delta(f) = (\text{id} \otimes \Delta)\Delta(f)$.

Again by Peter–Weyl theory, one obtains a vector space basis of $\mathcal{O}(G)$ consisting of the matrix coefficients u_{ij}^λ where λ ranges over $\text{Irr}(G)$ and $1 \leq i, j \leq \dim(\lambda)$. Here we abbreviate $\dim(\lambda) = \dim(\mathcal{H}_\lambda)$. If we write $\mathcal{O}(G)_\lambda$ for the linear span of the elements u_{ij}^λ for $1 \leq i, j \leq \dim(\lambda)$, we have a direct sum decomposition

$$\mathcal{O}(G) = \text{alg-} \bigoplus_{\lambda \in \text{Irr}(G)} \mathcal{O}(G)_\lambda.$$

Note that the coproduct of $\mathcal{O}(G)$ takes a particularly simple form in this picture; from the above formula for $\Delta(u_{ij}^\lambda)$ we see that it looks like the transpose of matrix multiplication.

Let u^λ be an irreducible unitary representation of G , and let u_{ij}^λ be the corresponding matrix elements in some fixed basis. The contragredient representation u^{λ^c} is given by the matrix $(u^{\lambda^c})_{ij} = S(u_{ji}^\lambda)$ where S is the antipode of $\mathcal{O}(G)$. In general u^{λ^c} is not unitary, but as any finite dimensional representation of G , it is unitarizable. The representations u^λ and $u^{\lambda^{cc}}$ are equivalent, and there exists a unique positive invertible intertwiner $F_\lambda : \mathcal{H}_\lambda \rightarrow \mathcal{H}_{\lambda^{cc}}$ satisfying $\text{tr}(F_\lambda) = \text{tr}(F_\lambda^{-1})$. The trace of F_λ is called the quantum dimension of \mathcal{H}_λ and denoted by $\dim_q(\lambda)$.

With this notation, the Schur orthogonality relations are

$$\varphi(u_{ij}^\lambda (u_{kl}^\eta)^*) = \delta_{\lambda\eta} \delta_{ik} \frac{1}{\dim_q(\lambda)} (F_\lambda)_{lj}$$

where $\lambda, \eta \in \text{Irr}(G)$ and $\varphi : C^r(G) \rightarrow \mathbb{C}$ is the Haar state of G . In the sequel we shall fix bases such that F_λ is a diagonal operator for all $\lambda \in \text{Irr}(G)$.

Remark 1.14. Let again G be a compact quantum group. While the comultiplication for the C^* -algebra $C^r(G)$ looks particularly simple in terms of matrix coefficients, dually the multiplication in the C^* -algebra $C^*(G)$ is easy to describe. More precisely, we have

$$C^*(G) \cong \bigoplus_{\lambda \in \text{Irr}(G)} \mathbb{K}(\mathcal{H}_\lambda),$$

where the right hand side denotes the c_0 -direct sum of the matrix algebras $\mathbb{K}(\mathcal{H}_\lambda)$. If u_{ij}^λ are the matrix coefficients in $\mathcal{O}(G)$, then the dual basis vectors ω_{ij}^λ , that is, the linear functionals on $\mathcal{O}(G)$ given by

$$\omega_{ij}^\lambda(u_{kl}^\eta) = \delta_{\lambda\eta} \delta_{ik} \delta_{jl}$$

form naturally a vector space basis of matrix units for the algebraic direct sum

$$\mathcal{D}(G) = \text{alg-} \bigoplus_{\lambda \in \text{Irr}(G)} \mathbb{K}(\mathcal{H}_\lambda)$$

inside $C_r^*(G)$.

Let us note that $\mathcal{D}(G)$ should not be confused with a quantum double, but there will be no conflicting notation in this regard appearing in this paper.

Let us also note that according to the Schur orthogonality relations, see [Remark 1.13](#), the functionals ω_{ij}^λ extend continuously to bounded linear functionals on $C^r(G)$.

1.2. Actions, crossed products and Takesaki–Takai duality

Let us now consider actions on C^* -algebras.

Definition 1.15. A (left) coaction of a Hopf C^* -algebra H on a C^* -algebra A is an injective nondegenerate $*$ -homomorphism $\alpha : A \rightarrow M(H \otimes A)$ such that the diagram

$$\begin{array}{ccc} A & \xrightarrow{\alpha} & M(H \otimes A) \\ \downarrow \alpha & & \downarrow \Delta \otimes \text{id} \\ M(H \otimes A) & \xrightarrow{\text{id} \otimes \alpha} & M(H \otimes H \otimes A) \end{array}$$

is commutative and $[\alpha(A)(H \otimes \mathbf{1})] = H \otimes A$.

If (A, α) and (B, β) are C^* -algebras equipped with coactions of H , then a $*$ -homomorphism $\varphi : A \rightarrow M(B)$ is called H -colinear if $\beta\varphi = (\text{id} \otimes \varphi)\alpha$.

An action of a locally compact quantum group G on a C^* -algebra A is a coaction $\alpha : A \rightarrow M(C_0^r(G) \otimes A)$ of $C_0^r(G)$ on A . We will also say that (A, α) is a G - C^* -algebra in this case. A $*$ -homomorphism $\varphi : A \rightarrow M(B)$ between G - C^* -algebras is called G -equivariant if it is $C_0^r(G)$ -colinear.

If H is a Hopf C^* -algebra and $\alpha : A \rightarrow M(H \otimes A)$ a coaction, then the density condition $[\alpha(A)(H \otimes \mathbf{1})] = H \otimes A$ implies in particular that the image of α is contained in the H -relative multiplier algebra of $H \otimes A$, defined by

$$M_H(H \otimes A) = \{m \in M(H \otimes A) \mid m(H \otimes \mathbf{1}), (H \otimes \mathbf{1})m \subset H \otimes A\}.$$

In contrast to the situation for ordinary multiplier algebras, the tensor product map $\text{id} \otimes \varphi : H \otimes A \rightarrow H \otimes B$ of a (possibly degenerate) $*$ -homomorphism $\varphi : A \rightarrow B$ admits a unique extension $M_H(H \otimes A) \rightarrow M_H(H \otimes B)$ to the relative multiplier algebra, which will again be denoted by $\text{id} \otimes \varphi$. If φ is injective, then this also holds for $\text{id} \otimes \varphi : M_H(H \otimes A) \rightarrow M_H(H \otimes B)$. We refer to [\[6, Appendix A\]](#) for further details on relative multiplier algebras.

Remark 1.16. For the construction of examples of Rokhlin actions we shall consider inductive limit actions of Hopf C^* -algebras. Assume H is a Hopf C^* -algebra and that $A_1 \rightarrow A_2 \rightarrow \cdots$ is an inductive system of H - C^* -algebras with coactions $\alpha_j : A_j \rightarrow M_H(H \otimes A_j)$ and injective equivariant connecting maps. Then the direct limit $A = \varinjlim A_j$ becomes an H - C^* -algebra in a canonical way. Firstly, we have $\varinjlim M_H(H \otimes A_j) \subset M_H(H \otimes A)$ naturally since our system $(A_j)_{j \in \mathbb{N}}$ has injective connecting maps. The maps α_j define a compatible family of $*$ -homomorphisms $\alpha_j : A_j \rightarrow M_H(H \otimes A_j) \rightarrow M_H(H \otimes A)$, and induce a $*$ -homomorphism $\alpha : A \rightarrow M_H(H \otimes A)$. Coassociativity of α and the density condition $[(H \otimes \mathbf{1})\alpha(A)] = H \otimes A$ follow from the corresponding properties of the coactions α_j . Injectivity of the map $\alpha : A \rightarrow M_H(H \otimes A)$ follows from injectivity of the maps α_j and of the connecting maps in the system $H \otimes A_j$.

Remark 1.17 (cf. [28]). Let G be a compact quantum group. Further below we will use the fact that any G - C^* -algebra A admits a spectral decomposition. In order to discuss this we review some further definitions and results.

Let G be a compact quantum group and let (A, α) be a G - C^* -algebra. Since G is compact, the coaction is an injective $*$ -homomorphism $\alpha : A \rightarrow C^r(G) \otimes A$ satisfying the coassociativity identity $(\Delta \otimes \text{id}) \circ \alpha = (\text{id} \otimes \alpha) \circ \alpha$ and the density condition $[(C^r(G) \otimes \mathbf{1})\alpha(A)] = C^r(G) \otimes A$. For $\lambda \in \text{Irr}(G)$ we let

$$A_\lambda = \{a \in A \mid \alpha(a) \in \mathcal{O}(G)_\lambda \odot A\}$$

be the λ -spectral subspace of A . Here we recall that $\mathcal{O}(G)_\lambda \subset \mathcal{O}(G)$ denotes the span of the matrix coefficients for \mathcal{H}_λ , see Remark 1.13.

The subspace A_λ is closed in A , and there is a projection operator $p_\lambda : A \rightarrow A_\lambda$ defined by

$$p_\lambda(a) = (\theta_\lambda \otimes \text{id})\alpha(a)$$

where

$$\theta_\lambda(x) = \dim_q(\lambda) \sum_{j=1}^{\dim(\lambda)} (F_\lambda)_{jj}^{-1} \varphi(x(u_{jj}^\lambda)^*).$$

By definition, the spectral subalgebra $\mathcal{S}(A) \subset A$ is the $*$ -subalgebra

$$\mathcal{S}(A) = \mathcal{S}_G(A) = \text{alg-} \bigoplus_{\lambda \in \text{Irr}(G)} A_\lambda.$$

From the Schur orthogonality relations and $[(C^r(G) \otimes \mathbf{1})\alpha(A)] = C^r(G) \otimes A$ it is easy to check that $\mathcal{S}(A)$ is dense in A .

For $\omega \in \mathcal{D}(G)$ and $a \in A$ let us define

$$a \cdot \omega = (\omega \otimes \text{id}) \circ \alpha(a).$$

Then the Schur orthogonality relations imply that $A \cdot \mathcal{D}(G)$, the linear span of all elements $a \cdot \omega$ as above, is equal to $\mathcal{S}(A)$. Moreover, from the coaction property of α it follows that A becomes a right $\mathcal{D}(G)$ -module in this way.

Definition 1.18. Let G be a locally compact quantum group and let A be a G - C^* algebra with coaction $\alpha : A \rightarrow M(C_0^r(G) \otimes A)$. The reduced crossed product $G \ltimes_{\alpha,r} A$ is the C^* -algebra

$$G \ltimes_{\alpha,r} A = [(C_0^r(\check{G}) \otimes \mathbf{1})\alpha(A)] \subset M(\mathbb{K}(L^2(G)) \otimes A).$$

Recall here that $C_0^r(\check{G}) = C_r^*(G)$ as a C^* -algebra, but equipped with the opposite comultiplication.

The reduced crossed product is equipped with a canonical dual action of \check{G} , which turns it into a \check{G} - C^* -algebra. More precisely, the dual action is given by comultiplication on the copy of $C_0^r(\check{G})$ and the trivial action on the copy of A inside $M(G \ltimes_{\alpha,r} A)$. By our second countability assumption, the crossed product $G \ltimes_{\alpha,r} A$ is separable provided A is separable.

For the purpose of reduced duality, we have to restrict ourselves to regular locally compact quantum groups [2]. All compact and discrete quantum groups are regular, so this is not an obstacle for the constructions we are interested in further below.

Remark 1.19. If G is a regular locally compact quantum group then the regular representation of $L^2(G)$ induces an action of G on the algebra $\mathbb{K}(L^2(G))$ of compact operators by conjugation. More generally, if A is any G - C^* -algebra we can turn the tensor product $\mathbb{K}(L^2(G)) \otimes A$ into a G - C^* -algebra by equipping $\mathbb{K}(L^2(G)) \otimes A \cong \mathbb{K}(L^2(G) \otimes A)$ with the conjugation action arising from the tensor product representation of G on the Hilbert A -module $\mathcal{E} = L^2(G) \otimes A$. Explicitly, following the notation in [26], we consider

$$\lambda(\xi \otimes a) = X_{12}^* \Sigma_{12} (\text{id} \otimes \alpha)(\xi \otimes a)$$

where $X = \Sigma V \Sigma$ and V is as in Remark 1.5. This determines a coaction $\lambda : \mathcal{E} \rightarrow M(C_0^r(G) \otimes \mathcal{E})$, which in turn corresponds to a unitary corepresentation

$$V_\lambda^* \in \mathbb{L}(C_0^r(G) \otimes \mathcal{E}) \cong M(C_0^r(G) \otimes \mathbb{K}(\mathcal{E})).$$

The conjugation action $\text{Ad}_\lambda : \mathbb{K}(\mathcal{E}) \rightarrow M(C_0^r(G) \otimes \mathbb{K}(\mathcal{E})) \cong \mathbb{L}(C_0^r(G) \otimes \mathcal{E})$ is then defined by

$$\text{Ad}_\lambda(T) = V_\lambda(\mathbf{1} \otimes T)V_\lambda^*,$$

and $\mathbb{K}(\mathcal{E})$ becomes a G - C^* -algebra in this way. Under the isomorphism $\mathbb{K}(L^2(G)) \otimes A \cong \mathbb{K}(L^2(G) \otimes A)$, this G -action is given by

$$\alpha_{\mathbb{K}}(T \otimes a) = \text{Ad}_{\lambda}(T \otimes a) = X_{12}^*(\mathbf{1} \otimes T \otimes \mathbf{1})\alpha(a)_{13}X_{12}, \quad T \in \mathbb{K}(L^2(G)), a \in A.$$

If G is a classical group then the resulting action is nothing but the diagonal action on the tensor product $\mathbb{K}(L^2(G)) \otimes A$. For further details we refer to [1,26].

Let us now state the Takesaki–Takai duality theorem for regular locally compact quantum groups [2], see [31, chapter 9] for a detailed exposition.

Theorem 1.20. *Let G be a regular locally compact quantum group and let A be a G - C^* -algebra. Then there is a natural isomorphism*

$$(\check{G} \ltimes_{\check{\alpha}, r} G \ltimes_{\alpha, r} A, \check{\alpha}) \cong (\mathbb{K}(L^2(G)) \otimes A, \alpha_{\mathbb{K}})$$

of G - C^* -algebras.

Let us give a brief sketch of the proof of Theorem 1.20 for the sake of convenience. Recall from Remark 1.5 that J is the modular conjugation of the left Haar weight of $L^\infty(G)$, and similarly \hat{J} the modular conjugation of the left Haar weight of $\mathcal{L}(G)$. We shall write $U = J\hat{J}$.

By definition, we have

$$\check{G} \ltimes_{\check{\alpha}, r} G \ltimes_{\alpha, r} A = [(UC_0^r(G)U \otimes \mathbf{1} \otimes \mathbf{1})(\hat{\Delta}^{\text{cop}}(C_r^*(G)) \otimes \mathbf{1})\alpha(A)_{23}].$$

Conjugating by W_{12}^* and using that $C_0^r(G)$ and $JC_0^r(G)J$ commute, this is isomorphic to

$$[(UC_0^r(G)U \otimes \mathbf{1} \otimes \mathbf{1})(C_r^*(G) \otimes \mathbf{1} \otimes \mathbf{1})(\text{id} \otimes \alpha)\alpha(A)] \cong [(UC_0^r(G)UC_r^*(G) \otimes \mathbf{1})\alpha(A)]$$

since α is injective. Moreover, $[UC_0^r(G)UC_r^*(G)]$ identifies with $\mathbb{K}(L^2(G))$. Hence the right hand side is isomorphic to $\mathbb{K}(L^2(G)) \otimes A$, taking into account that

$$[(\mathbb{K}(L^2(G)) \otimes \mathbf{1})\alpha(A)] = [(\mathbb{K}(L^2(G))C_0^r(G) \otimes \mathbf{1})\alpha(A)] = [\mathbb{K}(L^2(G))C_0^r(G) \otimes A]$$

by the density condition $[(C_0^r(G) \otimes \mathbf{1})\alpha(A)] = C_0^r(G) \otimes A$.

Let us also identify the bidual coaction. By construction, the bidual coaction $\check{\alpha}$ maps $UfU \otimes \mathbf{1} \in \mathbb{K}(L^2(G)) \otimes A$ for $f \in C_0^r(G)$ to $(\mathbf{1} \otimes U)\Delta(f)(\mathbf{1} \otimes U)$ and leaves $C_r^*(G) \otimes \mathbf{1}$ and $\alpha(A)$ fixed. Using the relations from Remark 1.5 one can show

$$\check{\alpha}(T \otimes a) = X_{12}^*(\mathbf{1} \otimes T \otimes \mathbf{1})\alpha(a)_{13}X_{12},$$

where $X = (\mathbf{1} \otimes U)W(\mathbf{1} \otimes U)$.

At some points we will also need the full crossed product $G \ltimes_{\alpha, f} A$ of a G - C^* -algebra (A, α) ; we refer to [26] for a review of its definition in terms of its universal property for covariant representations.

1.3. Braided tensor products

Finally, let us discuss Yetter–Drinfeld actions and braided tensor products. We refer to [26] for more information.

Definition 1.21. Let G be a locally compact quantum group. A G -YD- C^* -algebra is a C^* -algebra A together with a pair of actions $\alpha : A \rightarrow M(C_0^r(G) \otimes A)$ and $\gamma : A \rightarrow M(C_r^*(G) \otimes A)$ satisfying the Yetter–Drinfeld compatibility condition

$$(\sigma \otimes \text{id}) \circ (\text{id} \otimes \alpha) \circ \gamma = (\text{Ad}(W) \otimes \text{id}) \circ (\text{id} \otimes \gamma) \circ \alpha.$$

Here $W \in M(C_0^r(G) \otimes C_r^*(G))$ is the multiplicative unitary.

The braided tensor product, which we review next, generalizes at the same time the minimal tensor product of C^* -algebras and reduced crossed products. Roughly speaking, it allows one to construct a new C^* -algebra out of two constituent C^* -algebras, together with some prescribed commutation relations between the two factors.

Definition 1.22. Let G be a locally compact quantum group. Given a G -YD- C^* -algebra A and a G - C^* -algebra (B, β) , one defines the braided tensor product $A \boxtimes_G B = A \boxtimes B$ by

$$A \boxtimes B = [\gamma(A)_{12} \beta(B)_{13}] \subset M(\mathbb{K}(L^2(G)) \otimes A \otimes B).$$

The braided tensor product $A \boxtimes B$ becomes a G - C^* -algebra with coaction $\alpha \boxtimes \beta : A \boxtimes B \rightarrow M(C_0^r(G) \otimes A \boxtimes B)$ in such a way that the canonical embeddings $\iota_A : A \rightarrow M(A \boxtimes B)$ and $\iota_B : B \rightarrow M(A \boxtimes B)$ are G -equivariant. We write $a \boxtimes \mathbf{1}$ and $\mathbf{1} \boxtimes b$ for the images of elements $a \in A$ and $b \in B$ in $M(A \boxtimes B)$, respectively.

Example 1.23. A basic example of a G -YD-action is given by the C^* -algebra $A = C_0^r(G)$ for a regular locally compact quantum group G , equipped with $\alpha = \Delta$ and the adjoint action $\gamma(f) = \hat{W}^*(\mathbf{1} \otimes f)\hat{W}$.

We will mainly be interested in the special case of the braided tensor product construction where the first factor is equal to $C_0^r(G)$ with the Yetter–Drinfeld structure from Example 1.23.

Lemma 1.24. Let G be a regular locally compact quantum group. For any G - C^* -algebra A , there exists a G -equivariant $*$ -isomorphism

$$T_\alpha : (C_0^r(G) \boxtimes A, \Delta \boxtimes \alpha) \rightarrow (C_0^r(G) \otimes A, \Delta \otimes \text{id})$$

such that $T_\alpha(\mathbf{1} \boxtimes a) = \alpha(a)$ for all $a \in A \subset M(C_0^r(G) \boxtimes A)$ and $T_\alpha(f \boxtimes \mathbf{1}) = f \otimes \mathbf{1}$ for $f \in C_0^r(G) \subset M(C_0^r(G) \boxtimes A)$.

Proof. The map T_α is obtained from the identifications

$$\begin{aligned} C_0^r(G) \boxtimes A &= [\hat{W}_{12}^*(\mathbf{1} \otimes C_0^r(G) \otimes \mathbf{1})\hat{W}_{12}\alpha(A)_{13}] \\ &\cong [(\mathbf{1} \otimes C_0^r(G) \otimes \mathbf{1})\hat{W}_{12}\alpha(A)_{13}\hat{W}_{12}^*] \\ &\cong [(C_0^r(G) \otimes \mathbf{1} \otimes \mathbf{1})W_{12}^*\alpha(A)_{23}W_{12}] \\ &= [(C_0^r(G) \otimes \mathbf{1} \otimes \mathbf{1})(\text{id} \otimes \alpha)\alpha(A)] \\ &\cong [(C_0^r(G) \otimes \mathbf{1})\alpha(A)] \\ &= C_0^r(G) \otimes A, \end{aligned}$$

where we use $\hat{W} = \Sigma W^* \Sigma$ in the third step and the fact that α is injective in the penultimate step.

It is straightforward to check that $T_\alpha(a) = \alpha(a)$ for $a \in A \subset M(C_0^r(G) \boxtimes A)$ and $T_\alpha(f) = f \otimes \mathbf{1}$ for $f \in C_0^r(G) \subset M(C_0^r(G) \boxtimes A)$.

In particular, the canonical action of G on $C_0^r(G) \boxtimes A$ corresponds to the translation action on the first tensor factor in $C_0^r(G) \otimes A$ under this isomorphism. \square

2. Induced actions on sequence algebras

The theory of Rokhlin actions for compact quantum groups relies on the possibility of obtaining induced actions on the level of sequence algebras. In this section we shall recall a few facts on sequence algebras, and then discuss the construction of induced actions, separately for the case of discrete and compact quantum groups.

2.1. Sequence algebras

Let us first recall some notions related to sequence algebras, see [4] and [17]. If A is a C^* -algebra we write

$$\ell^\infty(A) = \{(a_n)_n \mid a_n \in A \text{ and } \sup_{n \in \mathbb{N}} \|a_n\| < \infty\}$$

for the C^* -algebra of bounded sequences with coefficients in A . Moreover denote by

$$c_0(A) = \{(a_n)_n \in \ell^\infty(A) \mid \lim_{n \rightarrow \infty} \|a_n\| = 0\} \subset \ell^\infty(A)$$

the closed two-sided ideal of sequences converging to zero.

Definition 2.1. Let A be a C^* -algebra. The sequence algebra A_∞ of A is

$$A_\infty = \ell^\infty(A)/c_0(A).$$

Given a bounded sequence $(a_n)_{n \in \mathbb{N}} \in \ell^\infty(A)$, the norm of the corresponding element in A_∞ is given by

$$\|[(a_n)_{n \in \mathbb{N}}]\| = \limsup_{n \rightarrow \infty} \|a_n\|.$$

Note moreover that A embeds canonically into A_∞ as (representatives of) constant sequences. We will frequently use this identification of A inside A_∞ in the sequel.

Notation 2.2. We denote by

$$\text{Ann}_\infty(A) = \{x \in A_\infty \mid xa = ax = 0 \text{ for all } a \in A\}$$

the two-sided annihilator of A inside A_∞ . Moreover, we write

$$D_{\infty,A} = [A \cdot A_\infty \cdot A] \subset A_\infty$$

for the hereditary subalgebra of A_∞ generated by A . Note that the embedding $A \hookrightarrow D_{\infty,A}$ is clearly nondegenerate. Finally, consider also the normalizer of $D_{\infty,A}$ inside A_∞ ,

$$\mathcal{N}(D_{\infty,A}, A_\infty) = \{x \in A_\infty \mid xD_{\infty,A} + D_{\infty,A}x \subset D_{\infty,A}\}.$$

We remark that $\text{Ann}_\infty(A)$ sits inside $\mathcal{N}(D_{\infty,A}, A_\infty)$ as a closed two-sided ideal.

The multiplier algebra of $D_{\infty,A}$ admits the following alternative description.

Proposition 2.3 (cf. [4, Proposition 1.5(1)] and [17, 1.9(4)]). *Let A be a σ -unital C^* -algebra. The canonical $*$ -homomorphism $\mathcal{N}(D_{\infty,A}, A_\infty) \rightarrow M(D_{\infty,A})$ given by the universal property of the multiplier algebra is surjective, and its kernel coincides with $\text{Ann}_\infty(A)$.*

Let us also note that the construction of $D_{\infty,A}$ is compatible with tensoring by the compacts.

Lemma 2.4 (cf. [4, 1.6]). *Let A be a C^* -algebra and let $\mathbb{K}(\mathcal{H})$ be the algebra of compact operators on a separable Hilbert space \mathcal{H} . The canonical embedding $\mathbb{K}(\mathcal{H}) \otimes A_\infty \rightarrow (\mathbb{K}(\mathcal{H}) \otimes A)_\infty$ induces an isomorphism $\mathbb{K}(\mathcal{H}) \otimes D_{\infty,A} \cong D_{\infty, \mathbb{K}(\mathcal{H}) \otimes A}$.*

Given a C^* -algebra equipped with an action of a quantum group G , we shall now discuss how to obtain induced actions on the sequence algebras introduced above.

2.2. Induced actions – discrete case

In the case of discrete quantum groups the situation is relatively simple. In fact, if G is a discrete quantum group then the C^* -algebra $C_0(G)$ of functions on G is a C^* -direct sum of matrix algebras. Explicitly, it is of the form

$$C_0(G) \cong \bigoplus_{\lambda \in \Lambda} \mathbb{K}(\mathcal{H}_\lambda)$$

where $\Lambda = \text{Irr}(\check{G})$ is the set of equivalence classes of irreducible representations of the dual compact quantum group, see [Remark 1.13](#).

If (A, α) is a G - C^* -algebra, then one of the defining conditions for the coaction $\alpha : A \rightarrow M(C_0(G) \otimes A)$ is that it factorizes over the $C_0(G)$ -relative multiplier algebra $M_{C_0(G)}(C_0(G) \otimes A)$. With the notation as above, we have

$$M_{C_0(G)}(C_0(G) \otimes A) \cong \prod_{\lambda \in \Lambda} \mathbb{K}(\mathcal{H}_\lambda) \otimes A,$$

that is, we can identify the relative multiplier algebra with the ℓ^∞ -product of the algebras $\mathbb{K}(\mathcal{H}_\lambda) \otimes A$. In other words, we have

$$\alpha : A \rightarrow \prod_{\lambda \in \Lambda} \mathbb{K}(\mathcal{H}_\lambda) \otimes A \subset M(C_0(G) \otimes A).$$

It follows that applying α componentwise induces a $*$ -homomorphism $\alpha^\infty : \ell^\infty(A) \rightarrow M(C_0(G) \otimes \ell^\infty(A))$ by considering the composition

$$\begin{aligned} \ell^\infty(A) &\rightarrow \prod_{n \in \mathbb{N}} \prod_{\lambda \in \Lambda} \mathbb{K}(\mathcal{H}_\lambda) \otimes A \\ &\cong \prod_{\lambda \in \Lambda} \prod_{n \in \mathbb{N}} \mathbb{K}(\mathcal{H}_\lambda) \otimes A \\ &\cong \prod_{\lambda \in \Lambda} \mathbb{K}(\mathcal{H}_\lambda) \otimes \prod_{n \in \mathbb{N}} A \subset M(C_0(G) \otimes \ell^\infty(A)), \end{aligned}$$

here we use that $\mathbb{K}(\mathcal{H}_\lambda)$ is finite dimensional for all $\lambda \in \Lambda$ in the second isomorphism.

Lemma 2.5. *Let G be a discrete quantum group and let (A, α) be a G - C^* -algebra. Then the map $\alpha^\infty : \ell^\infty(A) \rightarrow M(C_0(G) \otimes \ell^\infty(A))$ constructed above turns $\ell^\infty(A)$ into a G - C^* -algebra.*

Proof. Injectivity and coassociativity of α^∞ follow immediately from the corresponding properties of α . For the density condition

$$[(C_0(G) \otimes \mathbf{1})\alpha^\infty(\ell^\infty(A))] = C_0(G) \otimes \ell^\infty(A)$$

notice that it suffices to verify

$$[(pC_0(G) \otimes \mathbf{1})\alpha^\infty(\ell^\infty(A))] = pC_0(G) \otimes \ell^\infty(A)$$

for all finite rank (central) projections $p \in C_0(G)$. This in turn follows from the density condition for α , combined with fact that tensoring with finite dimensional algebras commutes with taking direct products. \square

The map α^∞ constructed above induces an injective $*$ -homomorphism $\alpha_\infty : A_\infty \rightarrow M(C_0(G) \otimes A_\infty)$ that fits into the following commutative diagram

$$\begin{array}{ccc} \ell^\infty(A) & \xrightarrow{\quad\quad\quad} & A_\infty \\ \downarrow \alpha^\infty & & \downarrow \alpha_\infty \\ M_{C_0(G)}(C_0(G) \otimes \ell^\infty(A)) & \longrightarrow & M_{C_0(G)}(C_0(G) \otimes A_\infty) \end{array}$$

Indeed, it suffices to observe that α^∞ maps $c_0(A)$ into $\prod_{\lambda \in \text{Irr}(G)} \mathbb{K}(\mathcal{H}_\lambda) \otimes c_0(A)$. Coassociativity and the density conditions for α_∞ are inherited from α^∞ . We therefore obtain the following result.

Lemma 2.6. *Let G be a discrete quantum group and let (A, α) be a G - C^* -algebra. Then the map $\alpha_\infty : A_\infty \rightarrow M(C_0(G) \otimes A_\infty)$ turns A_∞ into a G - C^* -algebra.*

2.3. Induced actions – compact case

Whereas for discrete quantum groups the extension of actions to sequence algebras always yields genuine actions, the situation for compact quantum groups is more subtle. Already classically, a strongly continuous action of a compact group G on a C^* -algebra A induces an action on $\ell^\infty(A)$ and A_∞ , but these induced actions typically fail to be strongly continuous, compare [4].

We shall address the corresponding problems in the quantum case by using an ad-hoc notion of equivariant $*$ -homomorphisms into sequence algebras. Our discussion also requires the technical assumption of coexactness.

Definition 2.7. A locally compact quantum group G is exact if the functor of taking reduced crossed products by G is exact. We say that G is coexact if the dual \hat{G} is exact.

It is well-known that a discrete quantum group G is exact if and only if $C_r^*(G)$ is an exact C^* -algebra, see [33, 1.28]. In other words, a compact quantum group G is coexact if and only if $C^*(G)$ is an exact C^* -algebra.

Remark 2.8. Let A and B be C^* -algebras. If B is exact, then there exists a canonical injective $*$ -homomorphism $B \otimes A_\infty \hookrightarrow (B \otimes A)_\infty$ coming from the following commutative diagram with exact rows

$$\begin{array}{ccccccc}
0 & \longrightarrow & B \otimes c_0(A) & \longrightarrow & B \otimes \ell^\infty(A) & \longrightarrow & B \otimes A_\infty \longrightarrow 0 \\
& & \downarrow \cong & & \downarrow & & \downarrow \\
0 & \longrightarrow & c_0(B \otimes A) & \longrightarrow & \ell^\infty(B \otimes A) & \longrightarrow & (B \otimes A)_\infty \longrightarrow 0
\end{array}$$

For similar reasons, we have a chain of natural inclusions $B \otimes M(D_{\infty,A}) \subset M(B \otimes D_{\infty,A}) \subset M(D_{\infty,B \otimes A})$ provided B is exact.

Now let G be a compact quantum group and let (A, α) be a G - C^* -algebra. The $*$ -homomorphism $\alpha : A \rightarrow C^r(G) \otimes A$ induces a $*$ -homomorphism

$$\alpha^\infty : \ell^\infty(A) \rightarrow \ell^\infty(C^r(G) \otimes A),$$

obtained by applying α componentwise, and also a $*$ -homomorphism

$$\alpha_\infty : A_\infty \rightarrow (C^r(G) \otimes A)_\infty.$$

The maps α^∞ and α_∞ , despite not being coactions in the sense of [Definition 1.15](#) in general, turn out to be good enough to obtain a tractable notion of equivariance and suitable crossed products, at least when G is coexact. The reason for this is [Remark 2.8](#), which is used in the definition below.

Definition 2.9. Let G be a coexact compact quantum group and let (A, α) and (B, β) be G - C^* -algebras. A $*$ -homomorphism $\varphi : A \rightarrow B_\infty$ is said to be G -equivariant if $\beta_\infty \circ \varphi = (\text{id} \otimes \varphi) \circ \alpha$, where both sides are viewed as maps from A into $(C^r(G) \otimes B)_\infty$. If φ is G -equivariant, then we also write $\varphi : (A, \alpha) \rightarrow (B_\infty, \beta_\infty)$.

Note in particular that if $\varphi : A \rightarrow B_\infty$ is a G -equivariant $*$ -homomorphism, then we automatically have $\beta_\infty \circ \varphi(A) \subset C^r(G) \otimes B_\infty$.

Remark 2.10. As indicated above, if G is a compact group and $\alpha : G \curvearrowright A$ is a strongly continuous action on a C^* -algebra, there always exists a (not necessarily strongly continuous) induced action of G on A_∞ . If (B, β) is another G -algebra, then it is easy to see that any $*$ -homomorphism $\varphi : A \rightarrow B_\infty$ is G -equivariant in the sense [2.9](#) if and only if it is G -equivariant in the usual sense.

Indeed, the equality $(\text{id} \otimes \varphi) \circ \alpha = \beta_\infty \circ \varphi$ clearly implies equivariance the usual sense. For the converse implication, one notes that if φ is equivariant in the usual sense, it maps A automatically into the continuous part of the action on B_∞ . Therefore, for $a \in A$ the equality of $(\text{id} \otimes \varphi) \circ \alpha(a) = \beta_\infty \circ \varphi(a)$ in $C(G) \otimes B_\infty = C(G, B_\infty)$ can be checked by evaluating both sides at the points of G .

In the quantum setting, we need a substitute of the continuous part of an action in order to define crossed products. We shall rely on the structure of compact quantum groups to obtain a construction suitable for the situation at hand.

Recall from Remark 1.13 that the dense Hopf $*$ -algebra $\mathcal{O}(G) \subset C^r(G)$ has a linear basis of elements of the form u_{ij}^λ where $\lambda \in \text{Irr}(G)$ and $1 \leq i, j, \leq \dim(\lambda)$. As explained in Remark 1.14, the linear functionals $\omega_{ij}^\lambda \in C^r(G)^*$ given by

$$\omega_{ij}^\lambda(u_{kl}^\eta) = \delta_{\lambda\eta} \delta_{ik} \delta_{jl}$$

span the space $\mathcal{D}(G)$, which can be viewed as the dense $*$ -subalgebra of $C_0(\hat{G})$ given by the algebraic direct sum of matrix algebras $\mathbb{K}(\mathcal{H}_\lambda)$ for $\lambda \in \text{Irr}(G)$. Moreover, in this picture the elements ω_{ij}^λ are matrix units in $\mathbb{K}(\mathcal{H}_\lambda)$, that is,

$$\omega_{ij}^\lambda \omega_{kl}^\eta = \delta_{\lambda\eta} \delta_{jk} \omega_{il}^\lambda$$

Let (A, α) be a G - C^* -algebra. Recall from Remark 1.17 that A becomes a right $\mathcal{D}(G)$ -module with the action

$$a \cdot \omega = (\omega \otimes \text{id}) \circ \alpha(a).$$

It is crucial for our purposes that such module structures also exist on $\ell^\infty(A)$ and A_∞ . Indeed, note that applying $\omega \otimes \text{id}$ in each component we obtain slice maps $\ell^\infty(C^r(G) \otimes A) \rightarrow \ell^\infty(A)$ and $(C^r(G) \otimes A)_\infty \rightarrow A_\infty$, which, by slight abuse of notation, will again be denoted by $\omega \otimes \text{id}$ in the sequel. We will also continue to use the notation $a \cdot \omega$ for the module structures obtained in this way.

In analogy with the constructions in Remark 1.17 we shall now define spectral subspaces of $\ell^\infty(A)$ and A_∞ , and use this to define corresponding continuous parts. Although the settings differ somewhat, this is similar to Kishimoto's definition of equicontinuous sequences for flows, cf. [20,19].

Definition 2.11. Let G be a coexact compact quantum group and let (A, α) be a G - C^* -algebra. The spectral subspaces of $\ell^\infty(A)$ and A_∞ with respect to α are defined by

$$\mathcal{S}(\ell^\infty(A)) = \ell^\infty(A) \cdot \mathcal{D}(G), \quad \mathcal{S}(A_\infty) = A_\infty \cdot \mathcal{D}(G),$$

respectively. The continuous parts of $\ell^\infty(A)$ and A_∞ with respect to α are defined by

$$\ell^{\infty, \alpha}(A) = [\mathcal{S}(\ell^\infty(A))] \subset \ell^\infty(A), \quad A_{\infty, \alpha} = [\mathcal{S}(A_\infty)] \subset A_\infty,$$

respectively.

At this point it is not immediately obvious that the subspaces in Definition 2.11 are closed under multiplication. We will show this further below.

Lemma 2.12. *The canonical map $\mathcal{S}(\ell^\infty(A)) \rightarrow \mathcal{S}(A_\infty)$ is surjective.*

Proof. Let $x \in \mathcal{S}(A_\infty)$. Then we can write $x = x \cdot p = (p \otimes \text{id})\alpha_\infty(x)$ for some finite rank idempotent $p \in \mathcal{D}(G)$. If $\tilde{x} \in \ell^\infty(A)$ is any lift of x , then $\tilde{x} \cdot p$ is a lift of x as well, which in addition is contained in $\mathcal{S}(\ell^\infty(A))$. \square

It follows from Lemma 2.12 that the canonical map $\ell^{\infty,\alpha}(A) \rightarrow A_{\infty,\alpha}$ is surjective.

Proposition 2.13. *Let G be a coexact compact quantum group and let (A, α) be a G - C^* -algebra. Then we have*

$$\begin{aligned}\mathcal{S}(\ell^\infty(A)) &= \{x \in \ell^\infty(A) \mid \alpha^\infty(x) \in \mathcal{O}(G) \odot \ell^\infty(A)\}; \\ \mathcal{S}(A_\infty) &= \{x \in A_\infty \mid \alpha_\infty(x) \in \mathcal{O}(G) \odot A_\infty\}.\end{aligned}$$

Moreover, both $\ell^{\infty,\alpha}(A)$ and $A_{\infty,\alpha}$ are G - C^* -algebras in a canonical way.

Proof. Let us consider first the assertions for $\mathcal{S}(\ell^\infty(A))$. By construction, for an element $x \in \mathcal{S}(\ell^\infty(A))$ satisfying $\alpha^\infty(x) \in \mathcal{O}(G) \odot \ell^\infty(A)$ we can write

$$\alpha^\infty(x) = \sum_{\lambda \in F} \sum_{i,j} u_{ij}^\lambda \otimes x_{ij}^\lambda$$

for some finite set $F \subset \text{Irr}(G)$ and elements $x_{ij}^\lambda \in \ell^\infty(A)$. By the definition of the counit $\epsilon : \mathcal{O}(G) \rightarrow \mathbb{C}$, we see that applying $\omega = \sum_{\lambda \in F} \sum_{i=1}^{\dim(\lambda)} \omega_{ii}^\lambda \in \mathcal{D}(G) \subset C^r(G)^*$ in the first tensor factor gives

$$x \cdot \omega = (\omega \otimes \text{id}) \circ \alpha^\infty(x) = (\epsilon \otimes \text{id}) \circ \alpha^\infty(x) = x.$$

This means $\{x \in \ell^\infty(A) \mid \alpha^\infty(x) \in \mathcal{O}(G) \odot \ell^\infty(A)\} \subset \mathcal{S}(\ell^\infty(A))$.

Conversely, write $x \in \mathcal{S}(\ell^\infty(A))$ as a finite sum $x = \sum_i y^i \cdot \omega^i$ of some elements $\omega^i \in \mathcal{D}(G)$ and $y^i \in \ell^\infty(A)$. We may assume without loss of generality that each ω^i is contained in $\mathbb{K}(\mathcal{H}_{\lambda_i})$ for some $\lambda_i \in \text{Irr}(G)$. Let $F \subset \text{Irr}(G)$ be the finite subset consisting of all λ_j . Writing $x = (x_n)_{n \in \mathbb{N}}$ and $y = (y_n)_{n \in \mathbb{N}}$ this means that each $x_n = \sum_i y_n^i \cdot \omega^i$ is contained in the subspace $\mathcal{S}(A)_F = A \cdot \mathcal{D}(G)_F \subset A \cdot \mathcal{D}(G)$, where $\mathcal{D}(G)_F = \sum_i \mathcal{D}(G)_{\lambda_i}$. Since $\mathcal{D}(G)_F$ is finite dimensional, we conclude

$$\alpha^\infty(x) \in \mathcal{O}(G) \odot \ell^\infty(A).$$

Here we use that the construction of ℓ^∞ -products is compatible with tensoring by finite dimensional spaces. We conclude $\mathcal{S}(\ell^\infty(A)) \subset \{x \in \ell^\infty(A) \mid \alpha^\infty(x) \in \mathcal{O}(G) \odot \ell^\infty(A)\}$.

The corresponding assertion for $\mathcal{S}(A_\infty)$ is obtained in a similar way. According to Lemma 2.12, we know that $x \in \mathcal{S}(A_\infty)$ is represented by an element $\tilde{x} \in \mathcal{S}(\ell^\infty(A))$, so the above argument shows $\alpha_\infty(x) \in \{z \in A_\infty \mid \alpha_\infty(z) \in \mathcal{O}(G) \odot A_\infty\}$ by construction of α_∞ . Conversely, if x satisfies $\alpha_\infty(x) \in \mathcal{O}(G) \odot A_\infty$ then the same argument as in the case $\ell^\infty(A)$ above shows $x \in \mathcal{S}(A_\infty)$.

As a consequence of these considerations we obtain in particular that $\mathcal{S}(\ell^\infty(A))$ and $\mathcal{S}(A_\infty)$ are $*$ -algebras, and hence $\ell^{\infty,\alpha}(A) \subset \ell^\infty(A)$ and $A_{\infty,\alpha} \subset A_\infty$ are C^* -subalgebras.

It remains to show that these C^* -algebras are G - C^* -algebras in a canonical way, with coactions induced by α^∞ and α_∞ , respectively.

Let us again first consider the case $\ell^{\infty,\alpha}(A)$. From coassociativity of α we obtain that α^∞ maps $\mathcal{S}(\ell^\infty(A))$ to $\mathcal{O}(G) \odot \mathcal{S}(\ell^\infty(A))$. Therefore it induces a $*$ -homomorphism $\ell^{\infty,\alpha}(A) \rightarrow C^r(G) \otimes \ell^{\infty,\alpha}(A) \subset C^r(G) \otimes \ell^\infty(A)$, which we will again denote by α^∞ . Injectivity of the latter map is clear. Similarly, the coaction identity $(\text{id} \otimes \alpha^\infty)\alpha^\infty = (\Delta \otimes \text{id})\alpha^\infty$ follows immediately from the coaction identity for α . For the density condition note that we can write $\mathbf{1} \otimes x = S(x_{(-2)})x_{(-1)} \otimes x_{(0)}$ for $x \in \mathcal{S}(\ell^\infty(A))$, using the Hopf algebra structure of $\mathcal{O}(G)$, and the Sweedler notation $\alpha^\infty(x) = x_{(-1)} \otimes x_{(0)}$. Hence

$$(\mathcal{O}(G) \odot \mathbf{1})\alpha^\infty(\mathcal{S}(\ell^\infty(A))) = \mathcal{O}(G) \odot \mathcal{S}(\ell^\infty(A)),$$

which implies $[(C^r(G) \otimes \mathbf{1})\alpha^\infty(\ell^{\infty,\alpha}(A))] = C^r(G) \otimes \alpha^\infty(\ell^{\infty,\alpha}(A))$ upon taking closures.

The case $A_{\infty,\alpha}$ is analogous. The considerations for $\mathcal{S}(\ell^\infty(A))$ above and [Lemma 2.12](#) imply that α_∞ maps $\mathcal{S}(A_\infty)$ to $\mathcal{O}(G) \odot \mathcal{S}(A_\infty)$. Therefore it induces a $*$ -homomorphism $A_{\infty,\alpha} \rightarrow C^r(G) \otimes A_{\infty,\alpha} \subset C^r(G) \otimes A_\infty$, which we will again denote by α_∞ . Coassociativity and density conditions are inherited from the corresponding properties of the coaction $\alpha^\infty : \ell^{\infty,\alpha}(A) \rightarrow C^r(G) \otimes \ell^{\infty,\alpha}(A)$. \square

Using [Proposition 2.13](#) we obtain an alternative way to describe the notion of equivariance introduced in [Definition 2.9](#)

Proposition 2.14. *Let G be a coexact compact quantum group and let (A, α) and (B, β) be G - C^* -algebras. For a $*$ -homomorphism $\varphi : A \rightarrow B_\infty$, the following are equivalent:*

- a) $\varphi : A \rightarrow B_\infty$ is G -equivariant;
- b) $\varphi(a \cdot \omega) = \varphi(a) \cdot \omega$ for all $\omega \in \mathcal{D}(G)$ and $a \in A$;
- c) $\varphi(A) \subset B_{\infty,\beta}$ and $\varphi : (A, \alpha) \rightarrow (B_{\infty,\beta}, \beta_\infty)$ is G -equivariant in the usual sense.

Proof. a) \Rightarrow b): Let $\omega \in \mathcal{D}(G)$ and $a \in A$, and recall $\mathcal{D}(G) \subset C^r(G)^*$. Applying $\omega \otimes \text{id}$ to both sides of the equality $(\text{id} \otimes \varphi) \circ \alpha(a) = \beta_\infty \circ \varphi(a)$, we obtain

$$\begin{aligned} \varphi(a \cdot \omega) &= (\text{id} \otimes \varphi)(\omega \otimes \text{id})\alpha(a) \\ &= (\omega \otimes \text{id})(\text{id} \otimes \varphi)\alpha(a) \\ &= (\omega \otimes \text{id})\beta_\infty \circ \varphi(a) \\ &= \varphi(a) \cdot \omega. \end{aligned}$$

b) \Rightarrow c) : As $\varphi(a \cdot \omega) = \varphi(a) \cdot \omega$ for all $\omega \in \mathcal{D}(G)$ and $a \in A$, it follows that $\varphi(\mathcal{S}(A)) \subset \mathcal{S}(B_\infty)$. Hence, $\beta_\infty \circ \varphi$ maps $\mathcal{S}(A)$ into $\mathcal{O}(G) \odot \mathcal{S}(B_\infty)$. For $a \in \mathcal{S}(A)$ and $\omega \in \mathcal{D}(G)$ we therefore compute

$$(\omega \otimes \text{id})\beta_\infty \varphi(a) = \varphi(a) \cdot \omega = \varphi(a \cdot \omega) = (\omega \otimes \text{id})(\text{id} \otimes \varphi)\alpha(a).$$

It is now straightforward to check that $\beta_\infty \circ \varphi(a) = (\text{id} \otimes \varphi) \circ \alpha(a)$ for all $a \in \mathcal{S}(A)$. As $\mathcal{S}(A) \subset A$ is dense, we conclude that $\varphi : A \rightarrow B_{\infty, \beta}$ is G -equivariant.

$c) \Rightarrow a)$: This implication follows immediately from the definitions. \square

We shall now define crossed products of induced actions on sequence algebras.

Definition 2.15. Let G be a coexact compact quantum group and let (A, α) be a G - C^* -algebra. We define

$$G \ltimes_{\alpha_\infty, r} A_\infty = G \ltimes_{\alpha_\infty, r} A_{\infty, \alpha},$$

that is, $G \ltimes_{\alpha_\infty, r} A_\infty$ is defined to be the crossed product of the continuous part of A_∞ with respect to the coaction $\alpha_\infty : A_{\infty, \alpha} \rightarrow C^r(G) \otimes A_{\infty, \alpha}$. In a similar way we define

$$G \ltimes_{\alpha_\infty, r} \ell^\infty(A) = G \ltimes_{\alpha_\infty, r} \ell^{\infty, \alpha}(A).$$

Remark 2.16. The notation introduced in Definition 2.15 will allow us to unify our exposition of several results in subsequent sections. Remark that the crossed products $G \ltimes_{\alpha_\infty, r} A_\infty$ and $G \ltimes_{\alpha_\infty, r} \ell^\infty(A)$ carry honest \check{G} - C^* -algebra structures given by the dual actions.

At a few points we will need a notion of equivariance for $*$ -homomorphisms with target $D_{\infty, B}$ or $M(D_{\infty, B})$.

Let G be a coexact compact quantum group and (B, β) be a G - C^* -algebra. Note that nondegeneracy of the $*$ -homomorphism $\beta : B \rightarrow C^r(G) \otimes B$ implies that

$$\beta_\infty(D_{\infty, B}) \subset D_{\infty, C^r(G) \otimes B} = (C^r(G) \otimes B)(C^r(G) \otimes B)_\infty(C^r(G) \otimes B)$$

is a nondegenerate C^* -subalgebra. Hence β_∞ induces a $*$ -homomorphism

$$M(D_{\infty, \beta}) \rightarrow M(D_{\infty, C^r(G) \otimes B}),$$

which we will again denote by β_∞ .

Definition 2.17. Let G be a coexact compact quantum group and let $(A, \alpha), (B, \beta)$ be G - C^* -algebras. A $*$ -homomorphism $\varphi : A \rightarrow M(D_{\infty, B})$ is called G -equivariant if $\beta_\infty \circ \varphi = (\text{id} \otimes \varphi) \circ \alpha$, where both sides are viewed as maps from A into $M(D_{\infty, C^r(G) \otimes B})$. If φ is G -equivariant, then we also write $\varphi : (A, \alpha) \rightarrow (M(D_{\infty, B}), \beta_\infty)$.

Remark 2.18. Note in particular that if $\varphi : A \rightarrow M(D_{\infty, B})$ is G -equivariant in the sense of Definition 2.17 then $\beta_\infty \circ \varphi(A) \subset C^r(G) \otimes M(D_{\infty, B}) \subset M(D_{\infty, C^r(G) \otimes B})$.

It is immediate from the definitions that a $*$ -homomorphism $\varphi : A \rightarrow D_{\infty,B}$ is G -equivariant as a $*$ -homomorphism $A \rightarrow B_{\infty}$ if and only if it is G -equivariant as a $*$ -homomorphism $A \rightarrow M(D_{\infty,B})$.

3. Equivariantly sequentially split $*$ -homomorphisms

In this section we discuss the notion of sequentially split $*$ -homomorphisms between G - C^* -algebras, which was studied in [4] in the case of actions by groups.

Definition 3.1 (cf. [4, 2.1, 3.3]). Let G be a quantum group which is either discrete or compact and coexact. Moreover let $(A, \alpha), (B, \beta)$ be G - C^* -algebras. We say that an equivariant $*$ -homomorphism $\varphi : (A, \alpha) \rightarrow (B, \beta)$ is equivariantly sequentially split if there exists a commutative diagram of G -equivariant $*$ -homomorphisms of the form

$$\begin{array}{ccc} (A, \alpha) & \xrightarrow{\quad} & (A_{\infty}, \alpha_{\infty}) \\ & \searrow \varphi & \nearrow \psi \\ & (B, \beta) & \end{array}$$

where the horizontal map is the standard embedding. If $\psi : (B, \beta) \rightarrow (A_{\infty}, \alpha_{\infty})$ is an equivariant $*$ -homomorphism fitting into the above diagram, then we say that ψ is an equivariant approximate left-inverse for φ .

An important feature of the theory of sequentially split $*$ -homomorphisms is that it is compatible with forming crossed product C^* -algebras. The proof makes use of the following fact.

Lemma 3.2. *Let G be a quantum group which is either discrete and exact or compact and coexact. Moreover let (A, α) be a G - C^* -algebra. Then there exists a \check{G} -equivariant $*$ -homomorphism $G \ltimes_{\alpha_{\infty},r} A_{\infty} \rightarrow (G \ltimes_{\alpha,r} A)_{\infty}$, compatible with the natural inclusions of $G \ltimes_{\alpha,r} A$ on both sides.*

Proof. Assume first that G is discrete and exact. Since taking reduced crossed products with G is exact, the canonical map $G \ltimes_{\alpha_{\infty},r} \ell^{\infty}(A) \rightarrow \ell^{\infty}(G \ltimes_{\alpha,r} A)$ induces a commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & G \ltimes_{c_0(\alpha),r} c_0(A) & \longrightarrow & G \ltimes_{\alpha_{\infty},r} \ell^{\infty}(A) & \longrightarrow & G \ltimes_{\alpha_{\infty},r} A_{\infty} \longrightarrow 0 \\ & & \downarrow \cong & & \downarrow & & \downarrow \\ 0 & \longrightarrow & c_0(G \ltimes_{\alpha,r} A) & \longrightarrow & \ell^{\infty}(G \ltimes_{\alpha,r} A) & \longrightarrow & (G \ltimes_{\alpha,r} A)_{\infty} \longrightarrow 0. \end{array}$$

Here $c_0(\alpha) : c_0(A) \rightarrow M_{C_0(G)}(C_0(G) \otimes c_0(A))$ denotes the restriction of α^{∞} to $c_0(A)$.

It is clear from the construction that the $*$ -homomorphism $G \ltimes_{\alpha_\infty, r} A_\infty \rightarrow (G \ltimes_{\alpha, r} A)_\infty$ is \check{G} -equivariant and compatible with the canonical inclusions of $G \ltimes_{\alpha, r} A$.

Assume now that G is compact and coexact. Let us abbreviate $\mathbb{K} = \mathbb{K}(L^2(G))$. Then the canonical map $\mathbb{K} \otimes \ell^\infty(A) \rightarrow \ell^\infty(\mathbb{K} \otimes A)$ induces the following commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{K} \otimes c_0(A) & \longrightarrow & \mathbb{K} \otimes \ell^\infty(A) & \longrightarrow & \mathbb{K} \otimes A_\infty \longrightarrow 0 \\ & & \downarrow \cong & & \downarrow & & \downarrow \\ 0 & \longrightarrow & c_0(\mathbb{K} \otimes A) & \longrightarrow & \ell^\infty(\mathbb{K} \otimes A) & \longrightarrow & (\mathbb{K} \otimes A)_\infty \longrightarrow 0 \end{array}$$

The middle vertical arrow restricts to a $*$ -homomorphism $G \ltimes_{\alpha_\infty, r} \ell^\infty(A) \hookrightarrow \ell^\infty(G \ltimes_{\alpha, r} A)$. By Lemma 2.12, the canonical \check{G} -equivariant map $G \rtimes_{\alpha_\infty, r} \ell^\infty(A) \rightarrow G \ltimes_{\alpha_\infty, r} A_\infty$ is surjective. Observe moreover that the canonical surjection $\ell^\infty(\mathbb{K} \otimes A) \rightarrow (\mathbb{K} \otimes A)_\infty$ restricts to the canonical surjection $\ell^\infty(G \ltimes_r A) \rightarrow (G \ltimes_r A)_\infty$. It follows that the embedding $\mathbb{K} \otimes A_\infty \hookrightarrow (\mathbb{K} \otimes A)_\infty$ restricts to an embedding $G \ltimes_{\alpha_\infty, r} A_\infty \hookrightarrow (G \ltimes_{\alpha, r} A)_\infty$. This map is clearly compatible with the canonical embeddings of $G \ltimes_{\alpha, r} A$. Moreover, as $G \ltimes_{\alpha_\infty, r} \ell^\infty(A) \rightarrow (G \ltimes_{\alpha, r} A)_\infty$ is \check{G} -equivariant, this also holds for $G \ltimes_{\alpha_\infty, r} A_\infty \rightarrow (G \ltimes_{\alpha, r} A)_\infty$. This finishes the proof. \square

Proposition 3.3. *Let G be quantum group which is either discrete and exact or compact and coexact. Moreover let (A, α) and (B, β) be G - C^* -algebras. Assume that $\varphi : (A, \alpha) \rightarrow (B, \beta)$ is an equivariantly sequentially split $*$ -homomorphism.*

Then the induced $$ -homomorphism $G \ltimes_r \varphi : G \ltimes_{\alpha, r} A \rightarrow G \ltimes_{\beta, r} B$ between the crossed products is \check{G} -equivariantly sequentially split.*

Proof. Let $\psi : (B, \beta) \rightarrow (A_\infty, \alpha_\infty)$ be an approximate left-inverse for φ . Passing to crossed products, we obtain a commutative diagram of \check{G} -equivariant $*$ -homomorphisms

$$\begin{array}{ccc} (G \ltimes_{\alpha, r} A, \check{\alpha}) & \xrightarrow{\quad \quad \quad} & (G \ltimes_{\alpha_\infty, r} A_\infty, \check{\gamma}) \\ & \searrow G \ltimes_r \varphi & \nearrow G \ltimes_r \psi \\ & (G \ltimes_{\beta, r} B, \check{\beta}) & \end{array}$$

where $\gamma = \alpha_\infty$. Composing $G \ltimes_r \psi$ with the \check{G} -equivariant $*$ -homomorphism $G \ltimes_{\alpha_\infty, r} A_\infty \rightarrow (G \ltimes_{\alpha, r} A)_\infty$ from Lemma 3.2 yields an equivariant approximate left-inverse for $G \ltimes_r \varphi$. \square

Let us next recall the definition of the fixed point algebra of an action of a compact quantum group.

Definition 3.4. Let G be a compact quantum group. For a G - C^* -algebra (A, α) we denote by

$$A^\alpha = \{a \in A \mid \alpha(a) = \mathbf{1} \otimes a\} \subset A$$

the C^* -subalgebra of fixed points.

Lemma 3.5. *Let G be a coexact compact quantum group and let (A, α) be a G - C^* -algebra. Then the canonical inclusion $(A^\alpha)_\infty \rightarrow (A_{\infty, \alpha})^{\alpha_\infty}$ is an isomorphism.*

Proof. Let $a \in (A_{\infty, \alpha})^{\alpha_\infty}$ be represented by $(a_n)_{n \in \mathbb{N}} \in \ell^\infty(A)$. Then

$$\lim_{n \rightarrow \infty} \|\alpha(a_n) - \mathbf{1} \otimes a_n\| = 0$$

by the fixed point condition. Applying the Haar state $\varphi : C^r(G) \rightarrow \mathbb{C}$ in the first tensor factor gives

$$\lim_{n \rightarrow \infty} \|(\varphi \otimes \text{id}) \circ \alpha(a_n) - a_n\| = 0$$

Since $(\varphi \otimes \text{id}) \circ \alpha$ maps A into A^α we conclude that a is represented by an element of $\ell^\infty(A^\alpha)$. \square

As we show next, a naturality property as in [Proposition 3.3](#) also holds for fixed point algebras of actions of compact quantum groups.

Proposition 3.6. *Let G be a coexact compact quantum group, and let $(A, \alpha), (B, \beta)$ be G - C^* -algebras. Assume that $\varphi : (A, \alpha) \rightarrow (B, \beta)$ is a G -equivariantly sequentially split $*$ -homomorphism.*

Then the induced $$ -homomorphism $\varphi : A^\alpha \rightarrow B^\beta$ is a sequentially split.*

Proof. Let $\psi : (B, \beta) \rightarrow (A_{\infty, \alpha})_{\infty}$ be an equivariant approximate left-inverse for φ . By equivariance of ψ , we have

$$\alpha_\infty \circ \psi(b) = (\text{id} \otimes \psi) \circ \beta(b) = \mathbf{1} \otimes \psi(b)$$

for all $b \in B^\beta$. Hence ψ maps B^β into $(A_{\infty, \alpha})^{\alpha_\infty}$. According to [Lemma 3.5](#) the latter identifies with $(A^\alpha)_\infty$, and therefore $\psi : B^\beta \rightarrow (A^\alpha)_\infty$ is an approximate left-inverse for $\varphi : A^\alpha \rightarrow B^\beta$. \square

The following stability result is an important feature for the theory of sequentially split $*$ -homomorphisms.

Proposition 3.7. *Let G be a quantum group which is either compact and coexact or discrete and exact. Moreover let $\varphi : (A, \alpha) \rightarrow (B, \beta)$ be a nondegenerate G -equivariant $*$ -homomorphism between G - C^* -algebras.*

Then φ is G -equivariantly sequentially split if and only if $\text{id} \otimes \varphi : (\mathbb{K}(L^2(G)) \otimes A, \alpha_{\mathbb{K}}) \rightarrow (\mathbb{K}(L^2(G)) \otimes B, \beta_{\mathbb{K}})$ is G -equivariantly sequentially split.

Proof. Let us first consider the case that G is compact and coexact.

Assume that φ is G -equivariantly sequentially split, and let $\psi : (B, \beta) \rightarrow (A_\infty, \alpha_\infty)$ be a G -equivariant approximate left inverse of φ . Then the map $\text{id} \otimes \psi : \mathbb{K}(L^2(G)) \otimes B \rightarrow \mathbb{K}(L^2(G)) \otimes A_\infty \subset (\mathbb{K}(L^2(G)) \otimes A)_\infty$ is G -equivariant, and yields an equivariant approximate left-inverse for $\text{id} \otimes \varphi$.

Conversely, assume that $\text{id} \otimes \varphi : (\mathbb{K}(L^2(G)) \otimes A, \alpha_\mathbb{K}) \rightarrow (\mathbb{K}(L^2(G)) \otimes B, \beta_\mathbb{K})$ is G -equivariantly sequentially split. Let $\Psi : (\mathbb{K}(L^2(G)) \otimes B, \beta_\mathbb{K}) \rightarrow ((\mathbb{K}(L^2(G)) \otimes A)_\infty, (\alpha_\mathbb{K})_\infty)$ be a G -equivariant approximate left-inverse. As φ is assumed to be non-degenerate, the image of Ψ is contained in $D_{\infty, \mathbb{K}(L^2(G)) \otimes A}$. Using the isomorphism $D_{\infty, \mathbb{K}(L^2(G)) \otimes A} \cong \mathbb{K}(L^2(G)) \otimes D_{\infty, A}$ from Lemma 2.4, we see that Ψ defines a non-degenerate $*$ -homomorphism from $\mathbb{K}(L^2(G)) \otimes B$ into $\mathbb{K}(L^2(G)) \otimes D_{\infty, A}$. Let us denote the extension $M(\mathbb{K}(L^2(G)) \otimes B) \rightarrow M(\mathbb{K}(L^2(G)) \otimes D_{\infty, A})$ to multiplier algebras again by Ψ .

We shall write Ψ_B for the restriction of Ψ to $B \cong \mathbf{1} \otimes B \subset M(\mathbb{K}(L^2(G)) \otimes B)$. Then $\Psi_B : B \rightarrow M(D_{\infty, \mathbb{K}(L^2(G)) \otimes A})$ is a $*$ -homomorphism whose image is contained in the relative commutant of $\mathbb{K}(L^2(G)) \otimes \mathbf{1}$. According to [4, 1.8], its image $\text{im}(\Psi_B)$ is therefore contained in $\mathbf{1} \otimes M(D_{\infty, A})$. Using again nondegeneracy of φ we see that $\text{im}(\Psi_B)$ is in fact contained in $\mathbf{1} \otimes D_{\infty, A}$. From these observations and the sequential split property we conclude that Ψ can be written in the form $\Psi = \text{id}_{\mathbb{K}(L^2(G))} \otimes \psi$ for a non-degenerate $*$ -homomorphism $\psi : B \rightarrow D_{\infty, A}$. It is easy to check that ψ is an approximate left-inverse for φ .

We claim that $\psi : B \rightarrow D_{\infty, A}$ is G -equivariant. For this consider a simple tensor $T \otimes b \in \mathbb{K}(L^2(G)) \otimes B$ and compute

$$\begin{aligned} (\text{id} \otimes \Psi)\beta_\mathbb{K}(T \otimes b) &= (\text{id} \otimes \Psi)(X_{12}^*(\mathbf{1} \otimes T \otimes \mathbf{1})\beta(b)_{13}X_{12}) \\ &= (\text{id} \otimes \text{id} \otimes \psi)(X_{12}^*(\mathbf{1} \otimes T \otimes \mathbf{1})\beta(b)_{13}X_{12}) \\ &= X_{12}^*(\mathbf{1} \otimes T \otimes \mathbf{1})((\text{id} \otimes \psi)\beta(b))_{13}X_{12} \end{aligned}$$

and

$$(\alpha_\mathbb{K})_\infty(\Psi(T \otimes b)) = (\alpha_\mathbb{K})_\infty(T \otimes \psi(b)) = X_{12}^*(\mathbf{1} \otimes T \otimes \mathbf{1})\alpha_\infty(\psi(b))_{13}X_{12}.$$

Here all expressions are viewed as elements of $(C^r(G) \otimes \mathbb{K}(L^2(G)) \otimes A)_\infty$. Equivariance of Ψ means that the above expressions are equal. We conclude $(\text{id} \otimes \psi)\beta(b) = \alpha_\infty(\psi(b))$ for all $b \in B$ as desired.

In the case that G is discrete and exact we can follow the above arguments almost word by word, in this case the situation is even slightly easier since all algebras involved are honest G - C^* -algebras. \square

Proposition 3.8. *Let G be a quantum group which is either compact and coexact or discrete and exact. Moreover assume that $(A, \alpha), (B, \beta)$ are separable G - C^* -algebras, and let $\varphi : (A, \alpha) \rightarrow (B, \beta)$ be a non-degenerate equivariant $*$ -homomorphism.*

Then φ is G -equivariantly sequentially split if and only if

$$\check{\varphi} = G \ltimes_r \varphi : (G \ltimes_{\alpha,r} A, \check{\alpha}) \rightarrow (G \ltimes_{\beta,r} B, \check{\beta})$$

is \check{G} -equivariantly sequentially split.

Proof. If φ is G -equivariantly sequentially split, then [Proposition 3.3](#) shows that $\check{\varphi}$ is \check{G} -equivariantly sequentially split. On the other hand, if $\check{\varphi}$ is \check{G} -equivariantly sequentially split, then [Proposition 3.3](#) implies that $\check{\check{\varphi}}$ is G -equivariantly sequentially split. Under the G -equivariant isomorphism given by Takesaki–Takai duality, see [Theorem 1.20](#), the map $\check{\check{\varphi}}$ corresponds to $\text{id}_{\mathbb{K}} \otimes \varphi : (\mathbb{K} \otimes A, \alpha_{\mathbb{K}}) \rightarrow (\mathbb{K} \otimes B, \beta_{\mathbb{K}})$. Here we abbreviate $\mathbb{K} = \mathbb{K}(L^2(G))$. Given an equivariant approximate left-inverse $(\check{G} \ltimes_{\check{\beta},r} G \ltimes_{\beta,r} B, \check{\check{\beta}}) \rightarrow ((\check{G} \ltimes_{\check{\alpha},r} G \ltimes_{\alpha,r} A)_{\infty}, \check{\check{\alpha}}_{\infty})$ for $\check{\check{\varphi}}$, componentwise application of Takesaki–Takai duality yields a G -equivariant approximate left-inverse $(\mathbb{K} \otimes B, \beta_{\mathbb{K}}) \rightarrow ((\mathbb{K} \otimes A)_{\infty}, (\alpha_{\mathbb{K}})_{\infty})$ for $\text{id}_{\mathbb{K}} \otimes \varphi$. Hence, $\text{id}_{\mathbb{K}} \otimes \varphi$ is G -equivariantly sequentially split. The claim now follows from [Proposition 3.7](#). \square

4. The Rokhlin property and approximate representability

In this section we introduce the key notions of this paper, namely the spatial Rokhlin property and spatial approximate representability for actions of quantum groups. Moreover we prove that these notions are dual to each other.

4.1. The spatial Rokhlin property

Let us start by defining the spatial Rokhlin property.

Definition 4.1. Let G be a coexact compact quantum group and (A, α) a separable G - C^* -algebra. We say that α has the spatial Rokhlin property if the second-factor embedding

$$\iota_{\alpha} = \mathbf{1} \boxtimes \text{id}_A : (A, \alpha) \rightarrow (C^*(G) \boxtimes A, \Delta \boxtimes \alpha)$$

is G -equivariantly sequentially split.

Remark 4.2. [Definition 4.1](#) is indeed a generalization of the classical notion of the Rokhlin property, see [\[4, 4.3\]](#). Indeed, for a classical compact group G , the braided tensor product $C(G) \boxtimes A$ agrees with the ordinary tensor product $C(G) \otimes A$. The term *spatial* in [Definition 4.1](#) refers to the fact that we have chosen to work with minimal (braided) tensor products; we will comment further on the implications of this choice in [Remark 4.9](#) below.

In special cases the Rokhlin property can be recast in the following way. Recall that if G is a compact quantum group and (A, α) a G - C^* -algebra we write $\mathcal{S}(A)$ for the spectral subalgebra of A . We shall use the Sweedler notation $\alpha(a) = a_{(-1)} \otimes a_{(0)}$ for the coaction $\alpha : \mathcal{S}(A) \rightarrow \mathcal{O}(G) \odot \mathcal{S}(A)$.

Proposition 4.3. *Let G be a coexact compact quantum group and (A, α) a separable G - C^* -algebra.*

a) *If α has the spatial Rokhlin property, then there exists a unital and G -equivariant $*$ -homomorphism $\kappa : (C^r(G), \Delta) \rightarrow (M(D_{\infty, A}), \alpha_{\infty})$ satisfying*

$$a\kappa(f) = \kappa(a_{(-2)}fS(a_{(-1)}))a_{(0)}$$

for all $f \in \mathcal{O}(G) \subset C^r(G)$ and $a \in \mathcal{S}(A) \subset A$. Moreover we have $\|\kappa(S(a_{(-1)}))a_{(0)}\| \leq \|a\|$ for all $a \in \mathcal{S}(A)$.

b) *Assume G is coamenable. If a $*$ -homomorphism $\kappa : C^r(G) \rightarrow M(D_{\infty, A})$ as in a) exists such that $\|\kappa(S(a_{(-1)}))a_{(0)}\| \leq \|a\|$ for all $a \in \mathcal{S}(A)$, then α has the spatial Rokhlin property.*

Proof. a) Assume first that α has the spatial Rokhlin property. Let

$$\psi : (C^r(G) \boxtimes A, \Delta \boxtimes \alpha) \rightarrow (A_{\infty}, \alpha_{\infty})$$

be an equivariant approximate left-inverse for ι_{α} as required by Definition 4.1. Since ι_{α} is non-degenerate, the image of this $*$ -homomorphism is contained in $D_{\infty, \alpha}$, and $\psi : C^r(G) \boxtimes A \rightarrow D_{\infty, A}$ is again nondegenerate. Let us also denote the unique strictly continuous extension of ψ to multipliers by the same letter, so that we have

$$\psi : M(C^r(G) \boxtimes A) \rightarrow M(D_{\infty, A}).$$

By equivariance of ψ , the unital $*$ -homomorphism

$$\kappa = \psi \circ \iota_{C^r(G)} : C^r(G) \rightarrow M(D_{\infty, A})$$

is also equivariant. According to the definition of the braided tensor product, we have

$$\iota_A(a)\iota_{C^r(G)}(f) = \iota_{C^r(G)}(a_{(-2)}fS(a_{(-1)}))\iota_A(a_{(0)})$$

for all $f \in \mathcal{O}(G)$ and $a \in \mathcal{S}(A)$. The desired twisted commutation relation for κ then follows by applying ψ . Moreover, the norm condition $\|\kappa(S(a_{(-1)}))a_{(0)}\| \leq \|a\|$ is a consequence of the formula $\kappa(S(a_{(-1)}))a_{(0)} = \psi(T_{\alpha}^{-1}(\mathbf{1} \otimes a))$ for $a \in \mathcal{S}(A)$ and the fact that ψ and the isomorphism T_{α}^{-1} from Lemma 1.24 are $*$ -homomorphisms between C^* -algebras.

b) Consider the map $\iota : \mathcal{S}(A) \rightarrow D_{\infty, A}$ given by $\iota(a) = \kappa(S(a_{(-1)}))a_{(0)}$. Then the commutation relation for κ gives

$$\iota(a)\kappa(f) = \kappa(S(a_{(-3)})a_{(-2)}fS(a_{(-1)}))a_{(0)} = \kappa(f)\iota(a)$$

for any $f \in \mathcal{O}(G)$, using the antipode relation for the Hopf algebra $\mathcal{O}(G)$. Using that S is antimultiplicative we therefore obtain

$$\begin{aligned}\iota(ab) &= \kappa(S(a_{(-1)}b_{(-1)}))a_{(0)}b_{(0)} \\ &= \kappa(S(b_{(-1)}))\kappa(S(a_{(-1)}))a_{(0)}b_{(0)} \\ &= \kappa(S(b_{(-1)}))\iota(a)b_{(0)} \\ &= \iota(a)\kappa(S(b_{(-1)}))b_{(0)} = \iota(a)\iota(b)\end{aligned}$$

for all $a, b \in \mathcal{S}(A)$. Moreover, using $S(h^*) = S^{-1}(h)^*$ for all $h \in \mathcal{O}(G)$ we have

$$\begin{aligned}\iota(a^*) &= \kappa(S(a_{(-1)}^*))a_{(0)}^* = \kappa(S^{-1}(a_{(-1)})^*)a_{(0)}^* \\ &= (a_{(0)}\kappa(S^{-1}(a_{(-1)})))^* \\ &= (\kappa(a_{(-2)}S^{-1}(a_{(-3)})S(a_{(-1)}))a_{(0)})^* \\ &= (\kappa(S(a_{(-1)}))a_{(0)})^* = \iota(a)^*\end{aligned}$$

for $a \in \mathcal{S}(A)$. It follows that ι is a $*$ -homomorphism. By assumption ι is bounded, so that it extends to a $*$ -homomorphism $\iota : A \rightarrow D_{\infty, A}$.

Combining κ and ι we obtain a $*$ -homomorphism $\psi = \kappa \otimes \iota : C(G) \otimes A \cong C(G) \otimes_{\max} A \rightarrow D_{\infty, A}$, using the universal property of the maximal tensor product and nuclearity of $C(G)$. Since κ is equivariant one checks that ι maps into the fixed point algebra of $A_{\infty, \alpha}$, and together with equivariance of κ it follows that $\psi : (C(G) \otimes A, \Delta \otimes \text{id}) \rightarrow (D_{\infty, A}, \alpha_{\infty})$ is G -equivariant. Using the isomorphism from [Lemma 1.24](#) we see that it defines an approximate left-inverse for $\iota_A : A \rightarrow C(G) \boxtimes A \cong C(G) \otimes A$. \square

Remark 4.4. Let us point out that the norm condition in part b) of [Proposition 4.3](#) is automatically satisfied if G is a finite quantum group. Indeed, in this case the spectral subalgebra $\mathcal{S}(A)$ is equal to A , and the claim follows from the fact that $*$ -homomorphisms between C^* -algebras are contractive. It can also be shown that the norm condition always holds if G is a classical compact group, but it seems unclear whether it is automatic in general.

Classically, a Rokhlin action of a compact group on an abelian C^* -algebra $C_0(X)$ induces a free action of G on X . In the quantum case, an analogue of the notion of freeness has been formulated by Ellwood in [\[7\]](#). Namely, an action $\alpha : A \rightarrow C^r(G) \otimes A$ of a compact quantum group G on a C^* -algebra A is called free if $[(1 \otimes A)\alpha(A)] = C^r(G) \otimes A$. It is shown in [\[7, Theorem 2.9\]](#) that this generalizes the classical concept of freeness.

Let us verify that the spatial Rokhlin property implies freeness also in the quantum case.

Proposition 4.5. *Let G be a coexact compact quantum group and let (A, α) be a separable G - C^* -algebra. If α has the spatial Rokhlin property, then it is free.*

Proof. Let $\psi : C^r(G) \boxtimes A \rightarrow A_\infty$ be an equivariant approximate left-inverse for the inclusion map $\iota_A : A \rightarrow C^r(G) \boxtimes A$. Notice that the action of G on $(C^r(G) \boxtimes A, \Delta \boxtimes \alpha) \cong (C^r(G) \otimes A, \Delta \otimes \text{id})$ is free, so that

$$[(\Delta \boxtimes \alpha)(C^r(G) \boxtimes A)(\mathbf{1} \otimes C^r(G) \boxtimes A)] = C^r(G) \otimes (C^r(G) \boxtimes A).$$

In fact, for any $f \in \mathcal{O}(G) \subset C^r(G)$ and $a \in A$ we find finitely many elements $x^i, y^i \in C^r(G) \boxtimes A$ such that $f \otimes (\mathbf{1} \boxtimes a) = \sum_i (\Delta \boxtimes \alpha)(x^i)(\mathbf{1} \otimes y^i)$ in $C^r(G) \otimes (C^r(G) \boxtimes A)$. Applying $\text{id} \otimes \psi$ to this equality and using equivariance, we obtain

$$f \otimes \psi(\mathbf{1} \boxtimes a) = f \otimes a = \sum_i \alpha_\infty \psi(x^i)(\mathbf{1} \otimes \psi(y^i))$$

in $(C^r(G) \otimes A)_\infty$. Consider lifts $(\tilde{x}_n^i)_{n \in \mathbb{N}}$ and $(\tilde{y}_n^i)_{n \in \mathbb{N}}$ in $\ell^\infty(A)$ for $\psi(x^i), \psi(y^i)$, respectively. Then

$$f \otimes a = \lim_{n \rightarrow \infty} \sum_i \alpha(\tilde{x}_n^i)(\mathbf{1} \otimes \tilde{y}_n^i),$$

which implies $C^r(G) \odot A \subset [\alpha(A)(\mathbf{1} \otimes A)]$, and hence also $C^r(G) \otimes A = [\alpha(A)(\mathbf{1} \otimes A)]$. \square

Let us point out that the Rokhlin property is strictly stronger than freeness; this is already the case classically. For instance, the antipodal action of $G = \mathbb{Z}_2$ on S^1 does not have the Rokhlin property.

Here comes the first main result of this paper:

Theorem 4.6. *Let G be a coexact compact quantum group and let (A, α) be a separable G - C^* -algebra. If α has the spatial Rokhlin property, then the two canonical embeddings*

$$A^\alpha \hookrightarrow A \quad \text{and} \quad G \ltimes_{\alpha, r} A \hookrightarrow \mathbb{K}(L^2(G)) \otimes A$$

are sequentially split. In particular, if A has any of the following properties, then so do the fixed-point algebra A^α and the crossed product $G \ltimes_{\alpha, r} A$:

- being simple;
- being nuclear and satisfying the UCT;
- having finite nuclear dimension or decomposition rank;
- absorbing a given strongly self-absorbing C^* -algebra \mathcal{D} .

Proof. Let $\psi : C^r(G) \boxtimes A \rightarrow A_\infty$ be an equivariant approximate left-inverse for the embedding $\iota_A : A \rightarrow C^r(G) \boxtimes A$. The resulting commutative diagram of equivariant $*$ -homomorphisms

$$\begin{array}{ccc} A & \xrightarrow{\quad} & A_\infty \\ & \searrow & \nearrow \psi \\ & C^r(G) \boxtimes A & \end{array}$$

induces a commutative diagram of $*$ -homomorphisms

$$\begin{array}{ccc} A^\alpha & \xrightarrow{\quad} & (A^\alpha)_\infty \\ & \searrow & \nearrow \psi \\ & A & \end{array}$$

by [Proposition 3.6](#). Notice here that [Lemma 1.24](#) implies that $(C^r(G) \boxtimes A)^{\Delta \boxtimes \alpha} \cong A$ in such a way that the canonical embeddings of A^α on both sides are compatible.

For the statement about the crossed product $G \ltimes_{\alpha,r} A$, observe that the spatial Rokhlin property for α means that $\alpha : A \rightarrow C^r(G) \otimes A$ is sequentially split, taking into account the isomorphism from [Lemma 1.24](#). According to [Proposition 3.3](#), it follows that the map

$$G \ltimes_r \alpha : G \ltimes_{\alpha,r} A \hookrightarrow G \ltimes_{\Delta \otimes \text{id}_A, r} (C^r(G) \otimes A) \cong (G \ltimes_{\Delta, r} C^r(G)) \otimes A$$

is sequentially split. Moreover, by the Takesaki–Takai duality [Theorem 1.20](#), we have $G \ltimes_{\Delta, r} C^r(G) \cong \mathbb{K}(L^2(G))$, and the resulting map $G \ltimes_{\alpha,r} A \rightarrow \mathbb{K}(L^2(G)) \otimes A$ is the standard embedding.

The asserted permanence properties are then a consequence of [\[4, Theorem 2.9\]](#). \square

4.2. Spatial approximate representability

Let us now define spatial approximate representability.

Let G be a discrete quantum group and (A, α) a separable G - C^* -algebra. Denote by $W_\alpha = (\text{id} \otimes \iota_G)(W) \in M(C_0(G) \otimes (G \ltimes_{\alpha,r} A))$ where $\iota_G : C_r^*(G) \rightarrow M(G \ltimes_{\alpha,r} A)$ is the canonical embedding. The unitary W_α implements the inner action of G on the crossed product, more precisely $\text{Ad}(W_\alpha^*)$ turns $M(G \ltimes_{\alpha,r} A)$ into a G - C^* -algebra such that

$$W_\alpha^*(1 \otimes a)W_\alpha = \alpha(a)$$

for all $a \in A \subset M(G \ltimes_{\alpha,r} A)$.

Definition 4.7. Let G be a discrete quantum group and (A, α) a separable G - C^* -algebra. We say that α is spatially approximately representable if the natural embedding

$$j_\alpha : (A, \alpha) \rightarrow (G \ltimes_{\alpha, r} A, \text{Ad}(W_\alpha^*))$$

is G -equivariantly sequentially split.

Proposition 4.8. *Let G be an exact discrete quantum group and (A, α) a separable G - C^* -algebra.*

- a) *If α is spatially approximately representable, then there exists a unitary representation $V \in M(C_0(G) \otimes D_{\infty, A})$ of G such that*

$$(\text{id} \otimes \alpha_\infty)(V) = V_{23}^* V_{13} V_{23}$$

and

$$V^*(\mathbf{1} \otimes a)V = \alpha(a)$$

for all $a \in A$.

- b) *If G is amenable and there exists a unitary representation $V \in M(C_0(G) \otimes D_{\infty, A})$ as in a), then α is spatially approximately representable.*

Proof. a) Assume that α is spatially approximately representable and let

$$\psi : G \ltimes_{\alpha, r} A \rightarrow A_\infty$$

be a G -equivariant approximate left-inverse for the embedding $A \rightarrow G \ltimes_{\alpha, r} A$. Since this embedding is nondegenerate, the image of ψ is contained in $D_{\infty, A}$, and $\psi : G \ltimes_{\alpha, r} A \rightarrow D_{\infty, A}$ is a nondegenerate $*$ -homomorphism. Let us denote the unique strictly continuous extension of ψ by the same letter, so that

$$\psi : M(G \ltimes_{\alpha, r} A) \rightarrow M(D_{\infty, A}).$$

We let

$$V = (\text{id} \otimes \psi)(W_\alpha) \in M(C_0(G) \otimes D_{\infty, A})$$

be the unitary representation corresponding to the restriction of ψ to $C_r^*(G)$. Equivariance of $\psi : G \ltimes_{\alpha, r} A \rightarrow D_{\infty, A}$ means

$$(\text{id} \otimes \psi)(W_\alpha^*(\mathbf{1} \otimes x)W_\alpha) = \alpha_\infty \circ \psi(x)$$

for all $x \in G \ltimes_{\alpha, r} A$. In particular, for every $a \in A$ we obtain

$$\alpha(a) = \alpha_\infty \circ \psi(a) = V^*(\mathbf{1} \otimes a)V.$$

Moreover, if $y = (\omega \otimes \text{id})(W_\alpha) \in M(G \rtimes_{\alpha,r} A)$ for $\omega \in \mathbb{L}(L^2(G))_*$, then

$$\begin{aligned} (\omega \otimes \text{id})(\text{id} \otimes \alpha_\infty)(V) &= \alpha_\infty \circ \psi(y) \\ &= (\text{id} \otimes \psi)(W_\alpha^*(\mathbf{1} \otimes y)W_\alpha) \\ &= (\omega \otimes \text{id} \otimes \text{id})(\text{id} \otimes \text{id} \otimes \psi)(W_{23}^*W_{13}W_{23}) \\ &= (\omega \otimes \text{id} \otimes \text{id})(V_{23}^*V_{13}V_{23}). \end{aligned}$$

Since this holds for all $\omega \in \mathbb{L}(L^2(G))_*$ we conclude $(\text{id} \otimes \alpha_\infty)(V) = V_{23}^*V_{13}V_{23}$.

b) Suppose that $V \in M(C_0(G) \otimes D_{\infty,A})$ is a unitary satisfying the conditions in a). Clearly, the canonical map $\iota : A \rightarrow D_{\infty,A}$ is equivariant and nondegenerate, and the formula

$$(\text{id} \otimes \iota) \circ \alpha(a) = V^*(\mathbf{1} \otimes \iota(a))V \quad \text{for all } a \in A$$

means that ι and V define a covariant pair. Hence they combine to a nondegenerate $*$ -homomorphism

$$\psi : G \rtimes_{\alpha,f} A \rightarrow M(D_{\infty,A})$$

such that $\psi \circ \iota(a) = a$ for all $a \in A$. Since G is amenable, we can identify the full crossed product $G \rtimes_{\alpha,f} A$ with the reduced crossed product $G \rtimes_{\alpha,r} A$. To verify that $\psi : G \rtimes_{\alpha,r} A \rightarrow M(D_{\infty,A})$ is G -equivariant, it suffices to check this separately on the copies of $C_r^*(G)$ and A inside $M(G \rtimes_{\alpha,r} A)$.

For $a \in A \subset G \rtimes_{\alpha,r} A$, the equivariance condition follows immediately from the relation $\psi \circ \iota(a) = a$. On $C_r^*(G)$ it is obtained by slicing the equation

$$\begin{aligned} (\text{id} \otimes \beta_\infty)(\text{id} \otimes \psi)(W_\beta) &= (\text{id} \otimes \beta_\infty)(V) \\ &= V_{23}^*V_{13}V_{23} \\ &= (\text{id} \otimes \text{id} \otimes \psi)(W_{23}^*W_{13}W_{23}), \end{aligned}$$

in the first tensor factor and using $C_r^*(G) = [(\mathbb{L}(L^2(G))_* \otimes \text{id})(W)]$. We conclude that ψ determines a G -equivariant approximate left-inverse for the inclusion $A \rightarrow G \rtimes_{\alpha,r} A$. \square

Remark 4.9. Definition 4.7 generalizes approximate representability for actions of discrete amenable groups, see [4, 4.23]. In the same way as already indicated in Remark 4.2, the term *spatial* in our definition is included since we work with minimal (braided) tensor products and reduced crossed products. In fact, approximate representability for classical discrete groups is defined in terms of the full crossed product instead, see [4, 4.23]. Notice that the trivial action of the free group \mathbb{F}_2 on \mathbb{C} is clearly approximately representable, but it is easily seen *not* to be spatially approximately representable in the sense of Definition 4.7.

It would therefore be more natural to develop the theory with maximal tensor products and full crossed products instead. However, this would mean in particular that one would have to work with full coactions taking values in maximal tensor products, which is technically less convenient.

Let us point out that all the above mentioned issues disappear for coamenable compact quantum groups and amenable discrete quantum groups, respectively; in these cases, we may omit the term *spatial*, and speak of the Rokhlin property and approximate representability.

4.3. Duality

We shall now show in several steps that the spatial Rokhlin property and spatial approximate representability are dual to each other.

Proposition 4.10. *Let G be a compact quantum group and (A, α) a separable G - C^* -algebra. Consider the G -equivariant $*$ -homomorphism*

$$\iota_\alpha = 1 \boxtimes \text{id}_A : (A, \alpha) \rightarrow (C^r(G) \boxtimes A, \Delta \boxtimes \alpha).$$

Then there exists a \check{G} -equivariant $$ -isomorphism*

$$\Psi_\alpha : (G \ltimes_{\Delta \boxtimes \alpha, r} (C^r(G) \boxtimes A), (\Delta \boxtimes \alpha)^\vee) \rightarrow (\check{G} \ltimes_{\check{\alpha}, r} (G \ltimes_{\alpha, r} A), \text{Ad}(W_\alpha^*))$$

that makes the following diagram commutative:

$$\begin{array}{ccc} (G \ltimes_{\alpha, r} A, \check{\alpha}) & \xrightarrow{G \ltimes_r \iota_\alpha} & (G \ltimes_{\Delta \boxtimes \alpha, r} (C^r(G) \boxtimes A), (\Delta \boxtimes \alpha)^\vee) \\ & \searrow j_{\check{\alpha}} & \downarrow \Psi_\alpha \\ & & (\check{G} \ltimes_{\check{\alpha}, r} (G \ltimes_{\alpha, r} A), \text{Ad}(W_\alpha^*)) \end{array}$$

Proof. Using [Lemma 1.24](#) and [Theorem 1.20](#), we obtain Ψ_α as the composition of the following identifications:

$$\begin{aligned} G \ltimes_{\Delta \boxtimes \alpha, r} (C^r(G) \boxtimes A) &\cong G \ltimes_{\Delta \otimes \text{id}_A, r} (C^r(G) \otimes A) \\ &\cong [(C_r^*(G) \otimes 1 \otimes 1) \Delta(C^r(G))_{12} (1 \otimes 1 \otimes A)] \\ &\cong [C_r^*(G) C^r(G) \otimes A] \\ &= \mathbb{K}(L^2(G)) \otimes A \\ &= [(UC^r(G) UC_r^*(G) \otimes 1) \alpha(A)] \\ &\cong \check{G} \ltimes_{\check{\alpha}, r} G \ltimes_{\alpha, r} A. \end{aligned}$$

Note that the copy of A inside $M(G \rtimes_{\Delta \boxtimes \alpha, r} (C^*(G) \boxtimes A))$ identifies with $\alpha(A) \subset M(\mathbb{K}(L^2(G)) \otimes A)$, and that the same holds for the copy of A inside $M(\check{G} \rtimes_{\check{\alpha}, r} G \rtimes_{\alpha, r} A)$. Similarly, the copies of $C_r^*(G)$ on both sides identify with $C_r^*(G) \otimes \mathbf{1} \subset M(\mathbb{K}(L^2(G)) \otimes A)$.

Moreover, the above identifications are compatible with the action of \check{G} on $\mathbb{K}(L^2(G)) \otimes A$ implemented by conjugation with $\Sigma \hat{V} \Sigma$. More precisely, the coaction

$$T \mapsto \text{Ad}(\Sigma_{12} \hat{V}_{12} \Sigma_{12})(\mathbf{1} \otimes T)$$

on $\mathbb{K}(L^2(G)) \otimes A$ corresponds to the dual coaction on $G \rtimes_{\Delta \boxtimes \alpha, r} (C^*(G) \boxtimes A)$ and to the conjugation coaction $\gamma = \text{Ad}(W_{\check{\alpha}}^*) : \check{G} \rtimes_{\hat{\alpha}, r} G \rtimes_{\alpha, r} A \rightarrow M(C_r^*(G)^{\text{cop}} \otimes \check{G} \rtimes_{\check{\alpha}, r} G \rtimes_{\alpha, r} A)$, given by $\gamma(T) = \hat{W}_{12}^U(\mathbf{1} \otimes T)(\hat{W}_{12}^U)^*_{12}$ where $W_{\check{\alpha}}^* = \hat{W}^U = (\mathbf{1} \otimes U)\hat{W}(\mathbf{1} \otimes U)$. For the latter observe $\Sigma \hat{V} \Sigma = (\mathbf{1} \otimes U)\hat{W}(\mathbf{1} \otimes U)$ and take into account the passage from $C_r^*(G)$ to $C_r^*(G)^{\text{cop}}$. \square

Proposition 4.11. *Let G be a discrete quantum group and (A, α) a separable G - C^* -algebra. Consider the G -equivariant inclusion*

$$j_{\alpha} : (A, \alpha) \rightarrow (G \rtimes_{\alpha, r} A, \text{Ad}(W_{\alpha}^*)).$$

Then there exists a \check{G} -equivariant $$ -isomorphism*

$$\Phi_{\alpha} : (G \rtimes_{\text{Ad}(W_{\alpha}^*), r} (G \rtimes_{\alpha, r} A), \text{Ad}(W_{\alpha}^*)^{\vee}) \rightarrow (C^*(\check{G}) \boxtimes (G \rtimes_{\alpha, r} A), \check{\Delta} \boxtimes \check{\alpha})$$

that makes the following diagram commutative:

$$\begin{array}{ccc} (G \rtimes_{\alpha, r} A, \check{\alpha}) & \xrightarrow{G \rtimes_r j_{\alpha}} & (G \rtimes_{\text{Ad}(W_{\alpha}^*), r} (G \rtimes_{\alpha, r} A), \text{Ad}(W_{\alpha}^*)^{\vee}) \\ & \searrow \iota_{\check{\alpha}} & \downarrow \Phi_{\alpha} \\ & & (C^*(\check{G}) \boxtimes (G \rtimes_{\alpha, r} A), \check{\Delta} \boxtimes \check{\alpha}) \end{array}$$

Proof. We obtain Φ_{α} as the composition of the following identifications:

$$\begin{aligned} G \rtimes_{\text{Ad}(W_{\alpha}^*)} (G \rtimes_{\alpha, r} A) &= [(C_r^*(G) \otimes \mathbf{1} \otimes \mathbf{1})W_{12}^*(\mathbf{1} \otimes C_r^*(G) \otimes \mathbf{1})\alpha(A)_{23}W_{12}] \\ &= [W_{12}^*(\hat{\Delta}^{\text{cop}}(C_r^*(G)) \otimes \mathbf{1})(\mathbf{1} \otimes C_r^*(G) \otimes \mathbf{1})\alpha(A)_{23}W_{12}] \\ &\cong [(\hat{\Delta}^{\text{cop}}(C_r^*(G)) \otimes \mathbf{1})(\mathbf{1} \otimes C_r^*(G) \otimes \mathbf{1})\alpha(A)_{23}] \\ &= [(C_r^*(G) \otimes C_r^*(G) \otimes \mathbf{1})\alpha(A)_{23}] \\ &\cong C_r^*(G)^{\text{cop}} \otimes (G \rtimes_{\alpha, r} A) \\ &= C^*(\check{G}) \otimes (G \rtimes_{\alpha, r} A) \\ &\cong C^*(\check{G}) \boxtimes (G \rtimes_{\alpha, r} A). \end{aligned}$$

Under these identifications, the copy of $C_r^*(G)$ inside $M(G \ltimes_{\alpha,r} A)$ on the left hand side gets identified with $\mathbf{1} \boxtimes (C_r^*(G) \otimes \mathbf{1})$ inside $C^r(\check{G}) \boxtimes (G \ltimes_{\alpha,r} A)$, and the copy of A in $G \ltimes_{\alpha,r} A$ is mapped to $\mathbf{1} \boxtimes \alpha(A)$. In other words, we indeed obtain a commutative diagram as desired.

Moreover, it is not hard to check that the dual action on $G \ltimes_{\text{Ad}(W_\alpha^*),r} (G \ltimes_{\alpha,r} A)$ corresponds to the action of $\hat{\Delta}^{\text{cop}} = \check{\Delta}$ on the first tensor factor of $C^r(\check{G}) \otimes (G \ltimes_{\alpha,r} A)$. It follows that Φ_α is \check{G} -equivariant. \square

As a consequence, we obtain the duality between the spatial Rokhlin property and spatial approximate representability.

Theorem 4.12. *Let G be a coexact compact quantum group and let (A, α) be a separable G - C^* -algebra. Then α has the spatial Rokhlin property if and only if $\check{\alpha}$ is spatially approximately representable.*

Dually, let G be an exact discrete quantum group and let (A, α) be a separable G - C^ -algebra. Then α is spatially approximately representable if and only if $\check{\alpha}$ has the spatial Rokhlin property.*

Proof. Let us first consider the case of compact quantum groups. By the general duality result from Proposition 3.8, we know that

$$\iota_\alpha : (A, \alpha) \hookrightarrow (C^r(G) \boxtimes A, \Delta \boxtimes \alpha)$$

is G -equivariantly sequentially split if and only if the induced $*$ -homomorphism

$$G \ltimes_r \iota_\alpha : (G \ltimes_{\alpha,r} A, \check{\alpha}) \rightarrow (G \ltimes_{\Delta \boxtimes \alpha,r} (C(G) \boxtimes A), (\Delta \boxtimes \alpha)^\vee)$$

is \check{G} -equivariantly sequentially split. By Proposition 4.10, there exists a commutative diagram of \check{G} -equivariant $*$ -homomorphisms

$$\begin{array}{ccc} (G \ltimes_{\alpha,r} A, \check{\alpha}) & \xrightarrow{G \ltimes_r \iota_\alpha} & (G \ltimes_{\Delta \boxtimes \alpha,r} (C(G) \boxtimes A), (\Delta \boxtimes \alpha)^\vee) \\ & \searrow j_{\check{\alpha}} & \downarrow \cong \\ & & (\check{G} \ltimes_{\check{\alpha},r} (G \ltimes_{\alpha,r} A), \text{Ad}(W_\alpha^*)) \end{array}$$

We conclude that ι_α is G -equivariantly sequentially split if and only if $j_{\check{\alpha}}$ is \check{G} -equivariantly sequentially split. This means that α has the spatial Rokhlin property if and only if $\check{\alpha}$ is spatially approximately representable.

The claim in the discrete case is proved in an analogous fashion. Again by Proposition 3.8, the G -equivariant $*$ -homomorphism

$$j_\alpha : (A, \alpha) \rightarrow (G \ltimes_{\alpha,r} A, \text{Ad}(W_\alpha^*))$$

is G -equivariantly sequentially split if and only if the induced $*$ -homomorphism

$$G \ltimes_r j_\alpha : (G \ltimes_{\alpha,r} A, \check{\alpha}) \rightarrow (G \ltimes_{\text{Ad}(W_\alpha^*),r} (G \ltimes_{\alpha,r} A), \text{Ad}(W_\alpha^*)^\vee)$$

is \check{G} -equivariantly sequentially split. By [Proposition 4.11](#), there exists a commutative diagram of \check{G} -equivariant $*$ -homomorphisms

$$\begin{array}{ccc} (G \ltimes_{\alpha,r} A, \check{\alpha}) & \xrightarrow{G \ltimes_r j_\alpha} & (G \ltimes_{\text{Ad}(W_\alpha^*),r} (G \ltimes_{\alpha,r} A), \text{Ad}(W_\alpha^*)^\vee) \\ & \searrow \iota_{\check{\alpha}} & \downarrow \cong \\ & & (C^r(\check{G}) \boxtimes (G \ltimes_{\alpha,r} A), \check{\Delta} \boxtimes \check{\alpha}) \end{array}$$

We conclude that j_α is G -equivariantly sequentially split if and only if $\iota_{\check{\alpha}}$ is \check{G} -equivariantly sequentially split. Hence α is spatially approximately representable if and only if $\check{\alpha}$ has the spatial Rokhlin property. \square

5. Rigidity of Rokhlin actions

In this section we provide a classification of actions of coexact compact quantum groups with the spatial Rokhlin property on separable C^* -algebras. This type of result was first obtained by Izumi in [\[15\]](#). Our basic approach follows Gardella–Santiago [\[11, Section 3\]](#), who proved corresponding results for finite group actions. We note that Gardella–Santiago have also announced the results of this section for actions of classical compact groups, see [\[10\]](#).

Recall that if (B, β) is a G - C^* -algebra for a compact quantum group G then $B^\beta \subset B$ denotes the fixed point subalgebra. We shall also write β for the induced coaction on the minimal unitarization \tilde{B} of B ; note that $\beta(\mathbf{1}) = \mathbf{1} \otimes \mathbf{1}$.

Definition 5.1. Let G be a compact quantum group. Let $\alpha : A \rightarrow C^r(G) \otimes A$ and $\beta : B \rightarrow C^r(G) \otimes B$ be two G -actions on C^* -algebras, and assume that A is separable. Let $\varphi_1, \varphi_2 : (A, \alpha) \rightarrow (B, \beta)$ be two equivariant $*$ -homomorphisms. We say that φ_1 and φ_2 are approximately G -unitarily equivalent, written $\varphi_1 \approx_{u,G} \varphi_2$, if there exists a sequence of unitaries $v_n \in \mathcal{U}(\tilde{B}^\beta)$ such that

$$\varphi_2(x) = \lim_{n \rightarrow \infty} v_n \varphi_1(x) v_n^* \quad \text{for all } x \in A.$$

Remark 5.2. For the trivial (quantum) group G , the above definition recovers the usual notion of approximate unitary equivalence between $*$ -homomorphisms. We write simply $\varphi_1 \approx_u \varphi_2$ instead of $\varphi_1 \approx_{u,G} \varphi_2$ in this case.

Proposition 5.3. Let G be a compact quantum group. Let $\alpha : A \rightarrow C^r(G) \otimes A$ and $\beta : B \rightarrow C^r(G) \otimes B$ be two G -actions on separable C^* -algebras. Let $\varphi : (A, \alpha) \rightarrow (B, \beta)$

and $\psi : (B, \beta) \rightarrow (A, \alpha)$ be two equivariant $*$ -homomorphisms such that $\psi \circ \varphi \approx_{u,G} \text{id}_A$ and $\varphi \circ \psi \approx_{u,G} \text{id}_B$. Then there exists an equivariant $*$ -isomorphism $\Phi : (A, \alpha) \rightarrow (B, \beta)$ such that $\Phi \approx_{u,G} \varphi$ and $\Phi^{-1} \approx_{u,G} \psi$.

Proof. This follows from a straightforward adaptation of the proof of [29, Corollary 2.3.4] to the setting of G -equivariant $*$ -homomorphism. For this, one requires the approximate intertwining from [29, Definition 2.3.1] to be (approximately) G -equivariant in the obvious way. The resulting $*$ -isomorphism $\Phi : A \rightarrow B$ then turns out to be α -to- β -equivariant. Moreover, the approximate unitary equivalences $\Phi \approx_u \varphi$ and $\Phi^{-1} \approx_u \psi$ that come out of the proof are indeed implemented by unitaries in $\mathcal{U}(\tilde{B}^\beta)$ and $\mathcal{U}(\tilde{A}^\alpha)$, respectively. \square

Let us now consider a series of partial results that will lead to the classification of Rokhlin actions.

Lemma 5.4. Let G be a compact quantum group. Let $\alpha : A \rightarrow C^*(G) \otimes A$ and $\beta : B \rightarrow C^*(G) \otimes B$ be two G -actions on separable C^* -algebras. Let $\varphi : A \rightarrow B$ be a $*$ -homomorphism that is equivariant modulo approximate unitary equivalence, i.e. $\beta \circ \varphi \approx_u (\text{id} \otimes \varphi) \circ \alpha$ as $*$ -homomorphisms between A and $C^*(G) \otimes B$. Then for every finite set $F \subset A$ and every $\varepsilon > 0$, there exists a unitary $v \in (C^*(G) \boxtimes B)^\sim$ such that

$$(\Delta \boxtimes \beta) \circ \text{Ad}(v) \circ (\mathbf{1} \boxtimes \varphi)(x) =_\varepsilon (\text{id} \otimes (\text{Ad}(v) \circ (\mathbf{1} \boxtimes \varphi))) \circ \alpha(x)$$

and

$$\|[(\mathbf{1} \boxtimes \varphi)(x), v]\| \leq \varepsilon + \|\beta \circ \varphi(x) - (\text{id} \otimes \varphi) \circ \alpha(x)\|$$

for all $x \in F$.

Proof. For convenience, the term id will always denote the identity map on $C^*(G)$ in this proof. Identity maps on other sets are decorated with the corresponding set.

Using our assumptions on φ , we may choose unitaries $u_n \in (C^*(G) \otimes B)^\sim$ such that

$$\text{Ad}(u_n) \circ \beta \circ \varphi \xrightarrow{n \rightarrow \infty} (\text{id} \otimes \varphi) \circ \alpha \quad (\text{e5.1})$$

in point-norm. By Lemma 1.24, we have the equivariant isomorphism

$$T_\beta : (C^*(G) \boxtimes B, \Delta \boxtimes \beta) \rightarrow (C^*(G) \otimes B, \Delta \otimes \text{id}_B) \quad (\text{e5.2})$$

satisfying

$$T_\beta(\mathbf{1} \boxtimes x) = \beta(x) \quad \text{for all } x \in B. \quad (\text{e5.3})$$

Let us also denote by T_β the obvious extension to the unitarizations. Set $v_n = T_\beta^{-1}(u_n)$. We calculate

$$\begin{aligned}
& \lim_{n \rightarrow \infty} (\Delta \boxtimes \beta) \circ \text{Ad}(v_n) \circ (\mathbf{1} \boxtimes \varphi) \\
&= \lim_{n \rightarrow \infty} (\Delta \boxtimes \beta) \circ \text{Ad}(T_\beta^{-1}(u_n)) \circ (\mathbf{1} \boxtimes \varphi) \\
&\stackrel{(e5.3)}{=} \lim_{n \rightarrow \infty} (\Delta \boxtimes \beta) \circ T_\beta^{-1} \circ \text{Ad}(u_n) \circ \beta \circ \varphi \\
&\stackrel{(e5.1)}{=} (\Delta \boxtimes \beta) \circ T_\beta^{-1} \circ (\text{id} \otimes \varphi) \circ \alpha \\
&\stackrel{(e5.2)}{=} (\text{id} \otimes T_\beta^{-1}) \circ (\Delta \otimes \text{id}_B) \circ (\text{id} \otimes \varphi) \circ \alpha \\
&= (\text{id} \otimes T_\beta^{-1}) \circ (\text{id} \otimes \text{id} \otimes \varphi) \circ (\Delta \otimes \text{id}_A) \circ \alpha \\
&= (\text{id} \otimes T_\beta^{-1}) \circ (\text{id} \otimes \text{id} \otimes \varphi) \circ (\text{id} \otimes \alpha) \circ \alpha \\
&= (\text{id} \otimes (T_\beta^{-1} \circ (\text{id} \otimes \varphi) \circ \alpha)) \circ \alpha \\
&\stackrel{(e5.1)}{=} \lim_{n \rightarrow \infty} (\text{id} \otimes (T_\beta^{-1} \circ \text{Ad}(u_n) \circ \beta \circ \varphi)) \circ \alpha \\
&= \lim_{n \rightarrow \infty} (\text{id} \otimes (\text{Ad}(v_n) \circ (\mathbf{1} \boxtimes \varphi))) \circ \alpha.
\end{aligned}$$

One should note that even though the existence of all these limits is a priori not clear at the beginning of this calculation, it follows a posteriori from the steps in this calculation.

Moreover, we calculate for all $x \in A$ that

$$\begin{aligned}
& \|[\mathbf{1} \boxtimes \varphi(x), v_n]\| \\
&= \|[(\mathbf{1} \boxtimes \varphi)(x), T_\beta^{-1}(u_n)]\| \\
&\stackrel{(e5.3)}{=} \|[(\beta \circ \varphi)(x), u_n]\| \\
&= \|(\text{Ad}(u_n) \circ \beta \circ \varphi)(x) - (\beta \circ \varphi)(x)\| \\
&\stackrel{(e5.1)}{\longrightarrow} \|(\text{id} \otimes \varphi) \circ \alpha(x) - \beta \circ \varphi(x)\|.
\end{aligned}$$

From these two calculations, it is clear that for given $F \subset A$ and $\varepsilon > 0$, any of the unitaries v_n satisfies the desired property for sufficiently large n . \square

Lemma 5.5. *Let G be a coexact compact quantum group. Let $\alpha : A \rightarrow C^*(G) \otimes A$ and $\beta : B \rightarrow C^*(G) \otimes B$ be two G -actions on separable C^* -algebras. Assume that β has the spatial Rokhlin property. Let $\varphi : A \rightarrow B$ be a $*$ -homomorphism that is equivariant modulo approximate unitary equivalence, i.e. $\beta \circ \varphi \approx_u (\text{id} \otimes \varphi) \circ \alpha$ as $*$ -homomorphisms between A and $C^*(G) \otimes B$. Then for every finite set $F \subset A$ and $\varepsilon > 0$, there exists a unitary $v \in \tilde{B}$ such that*

$$\beta \circ \text{Ad}(v) \circ \varphi(x) =_\varepsilon (\text{id} \otimes (\text{Ad}(v) \circ \varphi)) \circ \alpha(x)$$

and

$$\|[\varphi(x), v]\| \leq \varepsilon + \|\beta \circ \varphi(x) - (\mathbf{1} \otimes \varphi) \circ \alpha(x)\|$$

for all $x \in F$.

Proof. As β is assumed to have the spatial Rokhlin property, let

$$\psi : (C^*(G) \boxtimes B, \Delta \boxtimes \beta) \rightarrow (B_\infty, \beta_\infty)$$

be an equivariant $*$ -homomorphism satisfying

$$\psi(\mathbf{1} \boxtimes b) = b \quad \text{for all } b \in B. \quad (\text{e5.4})$$

We also denote by ψ the canonical extensions to the smallest unitarizations on both sides. Now let $F \subset A$ and $\varepsilon > 0$ be given. Apply [Lemma 5.4](#) and choose a unitary $w \in (C^*(G) \boxtimes B)^\sim$ such that

$$(\Delta \boxtimes \beta) \circ \text{Ad}(w) \circ (\mathbf{1} \boxtimes \varphi)(x) =_{\varepsilon/2} (\text{id} \otimes (\text{Ad}(w) \circ (\mathbf{1} \boxtimes \varphi))) \circ \alpha(x) \quad (\text{e5.5})$$

and

$$\|[(\mathbf{1} \boxtimes \varphi)(x), w]\| \leq \varepsilon/2 + \|\beta \circ \varphi(x) - (\text{id} \otimes \varphi) \circ \alpha(x)\| \quad (\text{e5.6})$$

for all $x \in F$. Set $v = \psi(w) \in (B_\infty)^\sim \subset \tilde{B}_\infty$. Combining the equivariance of ψ with [\(e5.4\)](#), [\(e5.5\)](#) and [\(e5.6\)](#), we obtain

$$\beta_\infty \circ \text{Ad}(v) \circ \varphi(x) =_{\varepsilon/2} (\text{id} \otimes (\text{Ad}(v) \circ \varphi)) \circ \alpha(x)$$

and

$$\|[\varphi(x), v]\| \leq \varepsilon/2 + \|\beta \circ \varphi(x) - (\text{id} \otimes \varphi) \circ \alpha(x)\|$$

for all $x \in F$. Now represent v by some sequence of unitaries $v_n \in \tilde{B}$. Then these equations translate to the conditions

$$\limsup_{n \rightarrow \infty} \|\beta \circ \text{Ad}(v_n) \circ \varphi(x) - (\text{id} \otimes (\text{Ad}(v_n) \circ \varphi)) \circ \alpha(x)\| \leq \varepsilon/2$$

and

$$\limsup_{n \rightarrow \infty} \|[\varphi(x), v_n]\| \leq \varepsilon/2 + \|\beta \circ \varphi(x) - (\text{id} \otimes \varphi) \circ \alpha(x)\|$$

for all $x \in F$. It follows that for sufficiently large n , any of the unitaries v_n satisfies the desired inequalities with respect to ε in place of $\varepsilon/2$. This finishes the proof. \square

Proposition 5.6 (cf. [\[11, 3.2\]](#)). *Let G be a coexact compact quantum group. Let $\alpha : A \rightarrow C^*(G) \otimes A$ and $\beta : B \rightarrow C^*(G) \otimes B$ be two G -actions on separable C^* -algebras. Assume that β has the spatial Rokhlin property. Let $\varphi : A \rightarrow B$ be a $*$ -homomorphism that is equivariant modulo approximate unitary equivalence, i.e. $\beta \circ \varphi \approx_u (\text{id} \otimes \varphi) \circ \alpha$ as $*$ -homomorphisms between A and $C^*(G) \otimes B$. Then there exists an equivariant $*$ -homomorphism $\psi : (A, \alpha) \rightarrow (B, \beta)$ with $\psi \approx_u \varphi$.*

Proof. Let

$$F_1 \subset F_2 \subset F_3 \subset \dots \subset A$$

be an increasing sequence of finite subsets with dense union. Let $(\varepsilon_n)_{n \in \mathbb{N}}$ be a decreasing sequence of strictly positive numbers with $\sum_{n=1}^{\infty} \varepsilon_n < \infty$. Using [Lemma 5.5](#) we find a unitary $v_1 \in \tilde{B}$ satisfying

$$\beta \circ \text{Ad}(v_1) \circ \varphi(x) =_{\varepsilon_1} (\text{id} \otimes (\text{Ad}(v_1) \circ \varphi)) \circ \alpha(x)$$

for all $x \in F_1$. Applying [Lemma 5.5](#) again (but now for $\text{Ad}(v_1) \circ \varphi$ in place of φ), we find a unitary $v_2 \in \tilde{B}$ satisfying

$$\beta \circ \text{Ad}(v_2 v_1) \circ \varphi(x) =_{\varepsilon_2} (\text{id} \otimes (\text{Ad}(v_2 v_1) \circ \varphi)) \circ \alpha(x)$$

and

$$\|[(\text{Ad}(v_1) \circ \varphi)(x), v_2]\| \leq \varepsilon_2 + \|\beta \circ \text{Ad}(v_1) \circ \varphi(x) - (\text{id} \otimes (\text{Ad}(v_1) \circ \varphi)) \circ \alpha(x)\|$$

for all $x \in F_2$. Applying [Lemma 5.5](#) again (but now for $\text{Ad}(v_2 v_1) \circ \varphi$ in place of φ), we find a unitary $v_3 \in \tilde{B}$ satisfying

$$\beta \circ \text{Ad}(v_3 v_2 v_1) \circ \varphi(x) =_{\varepsilon_3} (\text{id} \otimes (\text{Ad}(v_3 v_2 v_1) \circ \varphi)) \circ \alpha(x)$$

and

$$\|[(\text{Ad}(v_2 v_1) \circ \varphi)(x), v_3]\| \leq \varepsilon_3 + \|\beta \circ \text{Ad}(v_2 v_1) \circ \varphi(x) - (\mathbf{1} \otimes (\text{Ad}(v_2 v_1) \circ \varphi)) \circ \alpha(x)\|$$

for all $x \in F_3$. We inductively repeat this process and obtain a sequence of unitaries $v_n \in \tilde{B}$ satisfying

$$\beta \circ \text{Ad}(v_n \cdots v_1) \circ \varphi(x) =_{\varepsilon_n} (\text{id} \otimes (\text{Ad}(v_n \cdots v_1) \circ \varphi)) \circ \alpha(x) \quad (\text{e5.7})$$

for all $n \geq 1$ and

$$\begin{aligned} & \|[(\text{Ad}(v_{n-1} \cdots v_1) \circ \varphi)(x), v_n]\| \\ & \leq \varepsilon_n + \|\beta \circ \text{Ad}(v_{n-1} \cdots v_1) \circ \varphi(x) - (\text{id} \otimes (\text{Ad}(v_{n-1} \cdots v_n) \circ \varphi)) \circ \alpha(x)\| \end{aligned} \quad (\text{e5.8})$$

for all $x \in F_n$ and $n \geq 2$. For $m > n \geq k$ and $x \in F_k$ this implies

$$\begin{aligned}
& \| \operatorname{Ad}(v_m \cdots v_1) \circ \varphi(x) - \operatorname{Ad}(v_n \cdots v_1) \circ \varphi(x) \| \\
& \leq \sum_{j=n}^{m-1} \| \operatorname{Ad}(v_{j+1} \cdots v_1) \circ \varphi(x) - \operatorname{Ad}(v_j \cdots v_1) \circ \varphi(x) \| \\
& = \sum_{j=n}^{m-1} \| [(\operatorname{Ad}(v_j \cdots v_1) \circ \varphi)(x), v_{j+1}] \| \\
& \stackrel{(\text{e5.7}), (\text{e5.8})}{\leq} \sum_{j=n}^{m-1} \varepsilon_{j+1} + \varepsilon_j \leq 2 \cdot \sum_{j=n}^m \varepsilon_j.
\end{aligned}$$

As the ε_n were chosen as a 1-summable sequence and the union of the F_n is dense, this estimate implies that the sequence $\operatorname{Ad}(v_n \cdots v_1) \circ \varphi(x)$ is Cauchy for every $x \in A$. In particular, the point-norm limit $\psi = \lim_{n \rightarrow \infty} \operatorname{Ad}(v_n \cdots v_1) \circ \varphi$ exists and yields a well-defined $*$ -homomorphism from A to B . By construction we have $\psi \approx_u \varphi$, and the equivariance condition

$$\beta \circ \psi = (\operatorname{id} \otimes \psi) \circ \alpha$$

follows from (e5.7). This finishes the proof. \square

Lemma 5.7. *Let G be a compact quantum group. Let $\alpha : A \rightarrow C^r(G) \otimes A$ and $\beta : B \rightarrow C^r(G) \otimes B$ be two G -actions on separable C^* -algebras. Let $\varphi_1, \varphi_2 : (A, \alpha) \rightarrow (B, \beta)$ be two equivariant $*$ -homomorphisms. If $\varphi_1 \approx_u \varphi_2$, then $\mathbf{1} \boxtimes \varphi_1 \approx_{u,G} \mathbf{1} \boxtimes \varphi_2$ as equivariant $*$ -homomorphisms from A to $C^r(G) \boxtimes B$.*

Proof. Let $u_n \in \mathcal{U}(\tilde{B})$ be a sequence of unitaries satisfying

$$\operatorname{Ad}(u_n) \circ \varphi_1 \xrightarrow{n \rightarrow \infty} \varphi_2. \quad (\text{e5.9})$$

Using Lemma 1.24, we consider the equivariant isomorphism

$$T_\beta : (C^r(G) \boxtimes B, \Delta \boxtimes \beta) \rightarrow (C^r(G) \otimes B, \Delta \otimes \operatorname{id}_B)$$

that satisfies condition (e5.3). We shall also denote by T_β the obvious extension to the unitarizations. Set

$$v_n = T_\beta^{-1}(\mathbf{1} \otimes u_n) \in (C^r(G) \boxtimes B)^\sim.$$

As $\mathbf{1} \otimes u_n$ is in the fixed-point algebra of $\Delta \otimes \operatorname{id}_B$, it follows that v_n is in the fixed-point algebra of $\Delta \boxtimes \beta$. We have

$$\begin{aligned}
\operatorname{Ad}(v_n) \circ (\mathbf{1} \boxtimes \varphi_1) &\stackrel{(e5.3)}{=} T_\beta^{-1} \circ \operatorname{Ad}(\mathbf{1} \otimes u_n) \circ \beta \circ \varphi_1 \\
&= T_\beta^{-1} \circ \operatorname{Ad}(\mathbf{1} \otimes u_n) \circ (\mathbf{1} \otimes \varphi_1) \circ \alpha \\
&\stackrel{(e5.9)}{\longrightarrow} T_\beta^{-1} \circ (\mathbf{1} \otimes \varphi_2) \circ \alpha \\
&= T_\beta^{-1} \circ \beta \circ \varphi_2 \\
&\stackrel{(e5.3)}{=} \mathbf{1} \boxtimes \varphi_2.
\end{aligned}$$

This shows our claim. \square

Proposition 5.8. *Let G be a coexact compact quantum group. Let $\alpha : A \rightarrow C^*(G) \otimes A$, $\beta : B \rightarrow C^*(G) \otimes B$ and $\gamma : C \rightarrow C^*(G) \otimes C$ be three G -actions on separable C^* -algebras. Let*

$$\varphi_1, \varphi_2 : (A, \alpha) \rightarrow (B, \beta), \quad \psi : (B, \beta) \rightarrow (C, \gamma)$$

be equivariant $$ -homomorphisms. Assume that ψ is equivariantly sequentially split. Then $\varphi_1 \approx_{u,G} \varphi_2$ if and only if $\psi \circ \varphi_1 \approx_{u,G} \psi \circ \varphi_2$.*

Proof. If $\varphi_1 \approx_{u,G} \varphi_2$, then clearly $\psi \circ \varphi_1 \approx_{u,G} \psi \circ \varphi_2$. For the converse, assume that $\psi \circ \varphi_1 \approx_{u,G} \psi \circ \varphi_2$. Let $\kappa : (C, \gamma) \rightarrow (B_\infty, \beta_\infty)$ be an equivariant approximate left-inverse for ψ . Then $\kappa \circ \psi \circ \varphi_1 \approx_{u,G} \kappa \circ \psi \circ \varphi_2$, or in other words, $\varphi_1 \approx_{u,G} \varphi_2$ as equivariant $*$ -homomorphisms from A to $B_{\infty,\beta}$. Given $F \subset A$ and $\varepsilon > 0$, we therefore find some $u \in \mathcal{U}(\widetilde{(B_{\infty,\beta})}^{\beta_\infty})$ such that

$$\operatorname{Ad}(u) \circ \varphi_1(a) =_\varepsilon \varphi_2(a) \text{ for all } a \in F.$$

According to Lemma 3.5, we may choose a representing sequence $(u_n)_{n \in \mathbb{N}} \subset \mathcal{U}(\tilde{B}^\beta)$ for u . Picking a suitable member of this sequence, we find a unitary $v \in \mathcal{U}(\tilde{B}^\beta)$ such that

$$\operatorname{Ad}(v) \circ \varphi_1(a) =_{2\varepsilon} \varphi_2(a) \text{ for all } a \in F.$$

This shows that $\varphi_1 \approx_{u,G} \varphi_2$ as equivariant $*$ -homomorphisms from (A, α) to (B, β) . \square

Corollary 5.9 (cf. [11, 3.1]). *Let G be a coexact compact quantum group. Let $\alpha : A \rightarrow C^*(G) \otimes A$ and $\beta : B \rightarrow C^*(G) \otimes B$ be two G -actions on separable C^* -algebras. Assume that β has the spatial Rokhlin property. Let $\varphi_1, \varphi_2 : (A, \alpha) \rightarrow (B, \beta)$ be two equivariant $*$ -homomorphisms. Then $\varphi_1 \approx_u \varphi_2$ if and only if $\varphi_1 \approx_{u,G} \varphi_2$.*

Proof. We have to show that $\varphi_1 \approx_u \varphi_2$ implies $\varphi_1 \approx_{u,G} \varphi_2$. By Lemma 5.7, the amplified $*$ -homomorphisms are G -approximately unitarily equivalent, that is, $\mathbf{1} \boxtimes \varphi_1 \approx_{u,G} \mathbf{1} \boxtimes \varphi_2$. As β has the Rokhlin property, the canonical embedding $\mathbf{1} \boxtimes \operatorname{id}_B : (B, \beta) \rightarrow (C^*(G) \boxtimes B, \Delta \boxtimes \beta)$ is equivariantly sequentially split. Writing $\mathbf{1} \boxtimes \varphi_i = (\mathbf{1} \boxtimes \operatorname{id}_B) \circ \varphi_i$ for $i = 1, 2$, an application of Lemma 5.8 yields $\varphi_1 \approx_{u,G} \varphi_2$. This finishes the proof. \square

Here comes the main result of this section, which generalizes analogous results for finite group actions due to Izumi [15, 3.5], Nawata [25, 3.5] and Gardella–Santiago [11, 3.4]. It also generalizes the corresponding results for finite quantum groups by Osaka–Teruya [21, 10.7] and for classical compact groups by Gardella–Santiago [10].

Theorem 5.10. *Let G be a coexact compact quantum group. Let $\alpha, \beta : A \rightarrow C^*(G) \otimes A$ be two G -actions on a separable C^* -algebra. Assume that both have the spatial Rokhlin property. Then $\alpha \approx_u \beta$ as $*$ -homomorphisms if and only if there exists an equivariant isomorphism $\theta : (A, \alpha) \rightarrow (A, \beta)$ which is approximately inner as a $*$ -automorphism of A .*

Proof. First assume that $\theta : (A, \alpha) \rightarrow (A, \beta)$ is an equivariant $*$ -isomorphism which is approximately inner as a $*$ -automorphism. Then

$$\beta \approx_u \beta \circ \theta = (\text{id} \otimes \theta) \circ \alpha \approx_u \alpha.$$

Now assume that α and β are approximately unitarily equivalent. Then clearly

$$\beta \circ \text{id}_A = \beta \approx_u \alpha = (\text{id} \otimes \text{id}_A) \circ \alpha,$$

and analogously $\alpha \circ \text{id}_A \approx_u (\text{id} \otimes \text{id}_A) \circ \beta$. Since both α and β have the spatial Rokhlin property, it follows from Proposition 5.6 that there exist equivariant $*$ -homomorphisms $\varphi_1 : (A, \alpha) \rightarrow (A, \beta)$ and $\varphi_2 : (A, \beta) \rightarrow (A, \alpha)$ that are both approximately inner as $*$ -homomorphisms. Hence Corollary 5.9 implies $\varphi_1 \circ \varphi_2 \approx_{u,G} \text{id}_A$ and $\varphi_2 \circ \varphi_1 \approx_{u,G} \text{id}_A$. According to Proposition 5.3 we conclude that there exists an equivariant $*$ -isomorphism $\theta : (A, \alpha) \rightarrow (A, \beta)$ with $\theta \approx_{u,G} \varphi_1$. In particular, θ is also approximately inner as a $*$ -automorphism. \square

To conclude this section, we generalize the K -theory formula for fixed-point algebras of Rokhlin actions, which is originally due to Izumi and was recently extended by the first two authors.

Theorem 5.11 (cf. [15, 3.13] and [4, 4.9]). *Let G be a coexact compact quantum group. Let $\alpha : A \rightarrow C^*(G) \otimes A$ be an action on a separable C^* -algebra with the spatial Rokhlin property. Then the inclusion $A^\alpha \hookrightarrow A$ is injective in K -theory, and its image coincides with the subgroup*

$$K_*(A^\alpha) \cong \{x \in K_*(A) \mid K_*(\alpha)(x) = K_*(\mathbf{1} \otimes \text{id}_A)(x)\}.$$

Proof. If $x \in \text{im}(K_*(A^\alpha) \rightarrow K_*(A))$, then clearly $K_*(\alpha)(x) = K_*(\mathbf{1} \otimes \text{id}_A)(x)$. For the converse, let $x = [p] - [\mathbf{1}_k]$ be an element of $K_0(A)$, where $p \in M_n(\tilde{A})$ and $\mathbf{1}_k \in M_k(\tilde{A}) \subset M_n(\tilde{A})$ for some $k \leq n$ such that $p - \mathbf{1}_k \in M_n(A)$. Let us write

$$M_n(\tilde{\alpha}) : M_n(\tilde{A}) \rightarrow M_n((C^r(G) \otimes A)^\sim)$$

for the canonical extension of α to unitarizations and matrix amplifications.

Similarly, we write $M_n((\mathbf{1} \otimes \text{id}_A)^\sim)$ for the extension of $\mathbf{1} \otimes \text{id}_A$. If x satisfies $K_0(\alpha)(x) = K_0(\mathbf{1} \otimes \text{id}_A)(x)$ then

$$[M_n(\tilde{\alpha})(p)] - [M_n(\tilde{\alpha})(\mathbf{1}_k)] = [M_n((\mathbf{1} \otimes \text{id}_A)^\sim)(p)] - [M_n((\mathbf{1} \otimes \text{id}_A)^\sim)(\mathbf{1}_k)]$$

in $K_0(C^r(G) \otimes A)$. Notice that $M_n(\tilde{\alpha})(\mathbf{1}_k) = M_n((\mathbf{1} \otimes \text{id}_A)^\sim)(\mathbf{1}_k)$ by definition of $\tilde{\alpha}$, so that we get

$$[M_n(\tilde{\alpha})(p)] = [M_n((\mathbf{1} \otimes \text{id}_A)^\sim)(p)].$$

By definition of K_0 , we therefore find natural numbers m, l such that

$$M_n(\tilde{\alpha})(p) \oplus \mathbf{1}_m \oplus 0_l \sim_{\text{MvN}} M_n((\mathbf{1} \otimes \text{id}_A)^\sim)(p) \oplus \mathbf{1}_m \oplus 0_l$$

in $M_{n+m+l}((C^r(G) \otimes A)^\sim)$. Using the equivariant isomorphism

$$T = T_\alpha^{-1} : (C^r(G) \otimes A, \Delta \otimes \text{id}_A) \rightarrow (C^r(G) \boxtimes A, \Delta \boxtimes \alpha)$$

from [Lemma 1.24](#), we can view this as a relation in $M_{n+m+l}((C^r(G) \boxtimes A)^\sim)$. More precisely, using that $\alpha(a)$ and $\mathbf{1} \otimes a$ in $C^r(G) \otimes A$ for $a \in A$ correspond to the elements $\mathbf{1} \boxtimes a$ and $T(\mathbf{1} \otimes a)$ in $C(G) \boxtimes A$, respectively, we get

$$M_n((\mathbf{1} \boxtimes \text{id}_A)^\sim)(p) \oplus \mathbf{1}_m \oplus 0_l \sim_{\text{MvN}} M_n(\tilde{T}) \circ M_n((\mathbf{1} \otimes \text{id}_A)^\sim)(p) \oplus \mathbf{1}_m \oplus 0_l$$

in $M_{n+m+l}((C^r(G) \boxtimes A)^\sim)$. Write $n + m + l = r$. Since α has the Rokhlin property, let

$$\psi' : (C^r(G) \boxtimes A, \Delta \boxtimes \alpha) \rightarrow (A_\infty, \alpha_\infty)$$

be an equivariant $*$ -homomorphism with $\psi'(\mathbf{1} \boxtimes a) = a$ for all $a \in A$. We then consider

$$\psi = M_r(\psi') : M_r((C^r(G) \boxtimes A)^\sim) \rightarrow M_r(\tilde{A})_\infty,$$

which is an approximate left-inverse for $M_r((\mathbf{1} \boxtimes \text{id}_A)^\sim)$. Then

$$p \oplus \mathbf{1}_m \oplus 0_l \sim_{\text{MvN}} \psi \circ M_n(\tilde{T}) \circ M_n((\mathbf{1} \otimes \text{id}_A)^\sim)(p) \oplus \mathbf{1}_m \oplus 0_l$$

in $M_r(\tilde{A})_\infty$. Note that $M_n((\mathbf{1} \otimes \text{id}_A)^\sim)(p)$ is contained in the invariant part of $M_n((C^r(G) \otimes A)^\sim)$. That is,

$$M_n((\Delta \otimes \text{id}_A)^\sim) \circ M_n((\mathbf{1} \otimes \text{id}_A)^\sim)(p) = M_n((\mathbf{1} \otimes \mathbf{1} \otimes \text{id}_A)^\sim)(p)$$

in $M_n((C^r(G) \otimes C^r(G) \otimes A)^\sim)$. By equivariance of T and ψ , the same applies to

$$q_\infty = \psi \circ M_n(\tilde{T}) \circ M_n((\mathbf{1} \otimes \text{id}_A)^\sim)(p),$$

that is, the latter element satisfies

$$M_n(\tilde{\alpha}_\infty)(q_\infty) = M_n((\mathbf{1} \otimes \text{id}_{A_\infty})^\sim)(q_\infty).$$

Now the invariant part of $M_n(\tilde{A})_\infty$ equals $M_n((A^\alpha)^\sim)_\infty$ by Lemma 3.5. Since the relation of being a partial isometry with a fixed range projection is well-known to be weakly stable, this shows that there exists a projection $q \in M_r((A^\alpha)^\sim)$ such that

$$p \oplus \mathbf{1}_m \oplus 0_l \sim_{\text{MvN}} q$$

in $M_r(\tilde{A})$. Hence

$$x = [p] - [\mathbf{1}_k] = [p \oplus \mathbf{1}_m] - [\mathbf{1}_{m+k}] = [q] - [\mathbf{1}_{m+k}]$$

is contained in $\text{im}(K_0(A^\alpha) \rightarrow K_0(A))$ as desired.

For the statement about the K_1 -group, one uses suspension to reduce matters to K_0 , see the proof of [4, 4.9]. \square

6. Examples

In this final section we present some examples of Rokhlin actions.

Example 6.1. Let G be a coamenable compact quantum group acting on $A = C(G)$ by the regular coaction $\alpha = \Delta$. Then α has the spatial Rokhlin property.

Indeed, in this case the embedding $\iota_A : A \rightarrow C(G) \boxtimes A \cong C(G) \otimes A$ is given by $\iota_A = \Delta$. Since G is coamenable, the counit $\epsilon : \mathcal{O}(G) \rightarrow \mathbb{C}$ extends continuously to $C(G) = C^r(G)$, and $\text{id} \otimes \epsilon$ is an equivariant left-inverse for ι_A . Hence composition with the canonical embedding of $C(G) = A$ into A_∞ yields an equivariant approximate left-inverse.

The $*$ -homomorphism $\kappa : C(G) \rightarrow A_\infty$ corresponding to this Rokhlin action according to Proposition 4.3 is induced by the canonical embedding of $C(G)$ into its sequence algebra.

Remark 6.2. Let G be a finite quantum group and $\alpha : A \rightarrow C(G) \otimes A$ an action on a separable, unital C^* -algebra. In [21], Kodaka–Teruya introduce and study the Rokhlin property and approximate representability in this setting; in fact they also allow for twisted actions in their paper. It follows from Proposition 4.8 that α is spatially approximately representable in the sense of Definition 4.7 if and only if it is approximately representable in the sense of Kodaka–Teruya [21, Section 4]. As a consequence, Theorem 4.12 shows that α has the spatial Rokhlin property in the sense of Definition 4.1, if

and only if it has the Rokhlin property in the sense of Kodaka–Teruya [21, Section 5]. In particular, our definitions recover Kodaka–Teruya’s notions of the Rokhlin property and approximate representability and extend them to the non-unital setting. A substantial difference between our approach and [21] is that the duality of these two notions becomes a theorem rather than a definition.

Example 6.3. Let G be a finite quantum group of order $n = \dim(C(G))$. Then $B = M_n = M_n(\mathbb{C}) \cong \mathbb{K}(L^2(G))$ is a G -YD- C^* -algebra with the coactions $\beta : B \rightarrow C(G) \otimes B, \gamma : B \rightarrow C^*(G) \otimes B$ given by

$$\beta(T) = W^*(\mathbf{1} \otimes T)W, \quad \gamma(T) = \hat{W}^*(\mathbf{1} \otimes T)W^*,$$

respectively.

Let us write $B^{\boxtimes k} = B \boxtimes \cdots \boxtimes B$ for the k -fold braided tensor product. Note that the embeddings $M_n^{\boxtimes k} \rightarrow M_n \boxtimes M_n^{\boxtimes k} \cong M_n^{\boxtimes k+1}$ given by $T \mapsto \mathbf{1} \boxtimes T$ are G -equivariant. As explained in Remark 1.16, we may therefore form the inductive limit action $\alpha : A \rightarrow C(G) \otimes A$ of G on the corresponding inductive limit A .

We remark that A can be identified with the UHF-algebra M_{n^∞} . Indeed, the braided tensor product $M_n \boxtimes M_n$ is easily seen to be isomorphic to the ordinary tensor product $M_n \otimes M_n$ as a C^* -algebra, using that

$$\begin{aligned} M_n \boxtimes M_n &= [\hat{W}_{12}^*(\mathbf{1} \otimes M_n \otimes \mathbf{1})\hat{W}_{12}\beta(M_n)_{13}] \\ &\cong [(M_n \otimes \mathbf{1} \otimes \mathbf{1})(\text{id} \otimes \beta)\beta(M_n)] \\ &\cong [(M_n \otimes \mathbf{1})\beta(M_n)] \\ &\cong M_n \otimes M_n. \end{aligned}$$

An analogous statement holds for iterated braided tensor products.

We obtain a G -equivariant $*$ -homomorphism $\kappa : C(G) \rightarrow A_\infty$ by setting $\kappa(f) = [(\iota_k(f))_{k \in \mathbb{N}}]$, where $\iota_k : M_n \rightarrow A$ is the embedding into the k -th braided tensor factor of A . Moreover, for $a \in \iota_m(M_n) \subset A$ and $f \in C(G)$ we have

$$a\iota_k(f) = a_{(-2)}\iota_k(f)S(a_{(-1)})a_{(0)}$$

provided $k > m$. It follows that κ satisfies the commutation relations required by Proposition 4.3, and the norm condition in Proposition 4.3 b) is automatic since G is a finite quantum group. Hence (A, α) has the spatial Rokhlin property.

Proposition 6.4. *Let G be a coexact compact quantum group and \mathcal{D} a strongly self-absorbing C^* -algebra. Then there exists at most one conjugacy class of G -actions on \mathcal{D} with the spatial Rokhlin property.*

Proof. By [32, Corollary 1.12], any two unital $*$ -homomorphisms from \mathcal{D} to $C^*(G) \otimes \mathcal{D}$ are approximately unitarily equivalent. Therefore the claim follows from Theorem 5.10. \square

Remark 6.5. As a consequence of [Proposition 6.4](#) we see that the action of a finite quantum group G of order $n = \dim(C(G))$ constructed in [Example 6.3](#) is the unique Rokhlin action of G on M_{n^∞} up to conjugacy. In particular, it is conjugate to the action constructed by Kodaka–Teruya in [\[21, Section 7\]](#).

Finally, we shall construct a Rokhlin action of any coamenable compact quantum group on \mathcal{O}_2 . As a preparation, recall that an element a in a C^* -algebra A is called full if the closed two-sided ideal generated by a is equal to A .

Lemma 6.6. *Let G be a coamenable compact quantum group and let $\iota : C(G) \rightarrow \mathcal{O}_2$ be a unital embedding. Then $(\text{id} \otimes \iota) \circ \Delta(f)$ is a full element in $C(G) \otimes \mathcal{O}_2$ for any nonzero $f \in C(G)$.*

Proof. Let $f \in C(G) = C^r(G)$ be nonzero. To show that $h = (\text{id} \otimes \iota)(\Delta(f))$ is full it is enough to verify that $(\pi \otimes \text{id})(h)$ is nonzero for all irreducible representations π of $C(G)$. Indeed, if the ideal generated by h is proper, there must exist a primitive ideal of $C(G) \otimes \mathcal{O}_2$ containing h . Since \mathcal{O}_2 is nuclear and simple these ideals are of the form $I \otimes \mathcal{O}_2$ for primitive ideals $I \subset C(G)$, see [\[5, Theorem 3.3\]](#). Now if $\pi : C(G) \rightarrow \mathbb{L}(\mathcal{H}_\pi)$ is any $*$ -representation, then

$$(\pi \otimes \text{id})(\Delta(f)) = (\pi \otimes \text{id})(W)^*(\mathbf{1} \otimes f)(\pi \otimes \text{id})(W)$$

is nonzero in $\mathbb{L}(\mathcal{H}_\pi) \otimes C(G)$, and hence

$$(\text{id} \otimes \iota)((\pi \otimes \text{id})(\Delta(f))) = (\pi \otimes \iota)(\Delta(f)) = (\pi \otimes \text{id})(h)$$

is nonzero as well since ι is injective. \square

Theorem 6.7. *Let G be a coamenable compact quantum group. Then up to conjugacy, there exists a unique G -action on the Cuntz algebra \mathcal{O}_2 with the spatial Rokhlin property.*

Proof. According to [Proposition 6.4](#) it suffices to construct some G -action on \mathcal{O}_2 with the spatial Rokhlin property. Since $C^r(G) = C(G)$ is nuclear, hence in particular exact, there exists a unital embedding $\iota : C(G) \hookrightarrow \mathcal{O}_2$. For every $n \geq 0$, consider the unital $*$ -homomorphism $\Phi_n : C(G) \otimes \mathcal{O}_2^{\otimes n} \rightarrow C(G) \otimes \mathcal{O}_2^{\otimes n+1}$ given by

$$\Phi_n(x \otimes y) = ((\text{id} \otimes \iota) \circ \Delta)(x) \otimes y \quad \text{for all } x \in C(G), y \in \mathcal{O}_2^{\otimes n}.$$

Notice that $\Phi_n = \Phi_0 \otimes \text{id}_{\mathcal{O}_2^{\otimes n}}$ for all $n \geq 1$. We have

$$\begin{aligned}
(\Delta \otimes \text{id}_{\mathcal{O}_2}) \circ \Phi_0 &= (\Delta \otimes \text{id}_{\mathcal{O}_2}) \circ (\text{id} \otimes \iota) \circ \Delta \\
&= (\text{id} \otimes \text{id} \otimes \iota) \circ (\Delta \otimes \text{id}) \circ \Delta \\
&= (\text{id} \otimes \text{id} \otimes \iota) \circ (\text{id} \otimes \Delta) \circ \Delta \\
&= (\text{id} \otimes \Phi_0) \circ \Delta.
\end{aligned}$$

This means that Φ_0 is an injective equivariant $*$ -homomorphism

$$\Phi_0 : (C(G), \Delta) \rightarrow (C(G) \otimes \mathcal{O}_2, \Delta \otimes \text{id}_{\mathcal{O}_2}).$$

We thus also have that each

$$\Phi_n : (C(G) \otimes \mathcal{O}_2^{\otimes n}, \Delta \otimes \text{id}_{\mathcal{O}_2^{\otimes n}}) \rightarrow (C(G) \otimes \mathcal{O}_2^{\otimes n+1}, \Delta \otimes \text{id}_{\mathcal{O}_2^{\otimes n+1}})$$

is injective and equivariant. Define the inductive limit

$$(A, \alpha) = \varinjlim \left\{ (C(G) \otimes \mathcal{O}_2^{\otimes n}, \Delta \otimes \text{id}_{\mathcal{O}_2^{\otimes n}}), \Phi_n \right\},$$

where α denotes the inductive limit coaction, compare [Remark 1.16](#).

Notice that each building block in this inductive limit has the Rokhlin property, and moreover the first-factor embedding of $C(G)$ into $C(G) \otimes \mathcal{O}_2^{\otimes n}$ satisfies the required conditions from [Proposition 4.3 b\)](#) on the nose; see [Example 6.1](#). Similarly to what happens in [Example 6.3](#), the sequence

$$\kappa_n = \Phi_{n,\infty} \circ (\text{id} \otimes \mathbf{1}_{\mathcal{O}_2^{\otimes n}}) : C(G) \rightarrow A$$

yields a $*$ -homomorphism $\kappa : C(G) \rightarrow A_\infty$ satisfying the conditions in [Proposition 4.3 b\)](#). Hence α has the Rokhlin property.

By [Lemma 6.6](#), we know that Φ_0 , and thus also each Φ_n is a full $*$ -homomorphism. It follows that the inductive limit A is simple, and it is clearly separable, unital and nuclear. Moreover A is \mathcal{O}_2 -absorbing by [\[32, 3.4\]](#). This implies $A \cong \mathcal{O}_2$ due to Kirchberg–Phillips [\[18\]](#). \square

References

- [1] S. Baaj, G. Skandalis, C^* -algèbres de Hopf et théorie de Kasparov équivariante, *K-Theory* 2 (6) (1989) 683–721, <http://dx.doi.org/10.1007/BF00538428>.
- [2] S. Baaj, G. Skandalis, Unitaires multiplicatifs et dualité pour les produits croisés de C^* -algèbres, *Ann. Sci. Éc. Norm. Supér.* (4) 26 (4) (1993) 425–488, http://www.numdam.org/item?id=ASENS_1993_4_26_4_425_0.
- [3] S. Baaj, G. Skandalis, S. Vaes, Non-semi-regular quantum groups coming from number theory, *Comm. Math. Phys.* 235 (1) (2003) 139–167, <http://dx.doi.org/10.1007/s00220-002-0780-6>.
- [4] S. Barlak, G. Szabó, Sequentially split $*$ -homomorphisms between C^* -algebras, *Int. J. Math* 27 (12) (2016), 48 pp., <http://dx.doi.org/10.1142/S0129167X16501056>.

- [5] B.E. Blackadar, Infinite tensor products of C^* -algebras, *Pacific J. Math.* 72 (2) (1977) 313–334.
- [6] S. Echterhoff, S. Kaliszewski, J. Quigg, I. Raeburn, A Categorical Approach to Imprimitivity Theorems for C^* -Dynamical Systems, *Mem. Amer. Math. Soc.*, vol. 180, American Mathematical Society, 2006.
- [7] D.A. Ellwood, A new characterisation of principal actions, *J. Funct. Anal.* 173 (1) (2000) 49–60, <http://dx.doi.org/10.1006/jfan.2000.3561>.
- [8] E. Gardella, Compact group actions on C^* -algebras: classification, non-classifiability, and crossed products, and rigidity results for L^p -operator algebras, Ph.D. thesis, University of Oregon, 2015.
- [9] E. Gardella, Crossed products by compact group actions with the Rokhlin property, *J. Noncommut. Geom.* (2016), in press, <http://arxiv.org/abs/1408.1946>.
- [10] E. Gardella, L. Santiago, Classification of actions of compact groups with the Rokhlin property, 2016, in preparation.
- [11] E. Gardella, L. Santiago, Equivariant $*$ -homomorphisms, Rokhlin constraints and equivariant UHF-absorption, *J. Funct. Anal.* 270 (7) (2016) 2543–2590.
- [12] R. Herman, V. Jones, Period two automorphisms of UHF C^* -algebras, *J. Funct. Anal.* 45 (2) (1982) 169–176.
- [13] R. Herman, A. Ocneanu, Stability for integer actions on UHF C^* -algebras, *J. Funct. Anal.* 59 (1984) 132–144.
- [14] I. Hirshberg, W. Winter, Rokhlin actions and self-absorbing C^* -algebras, *Pacific J. Math.* 233 (1) (2007) 125–143.
- [15] M. Izumi, Finite group actions on C^* -algebras with the Rohlin property I, *Duke Math. J.* 122 (2) (2004) 233–280.
- [16] M. Izumi, Finite group actions on C^* -algebras with the Rohlin property II, *Adv. Math.* 184 (1) (2004) 119–160.
- [17] E. Kirchberg, Central sequences in C^* -algebras and strongly purely infinite algebras, in: *Operator Algebras: The Abel Symposium 2004*, vol. 1, Springer, Berlin, 2006, pp. 175–231.
- [18] E. Kirchberg, N.C. Phillips, Embedding of exact C^* -algebras in the Cuntz algebra \mathcal{O}_2 , *J. Reine Angew. Math.* 525 (2000) 17–53.
- [19] A. Kishimoto, Rohlin property for flows, in: *Advances in Quantum Dynamics*, South Hadley, MA, 2002, in: *Contemp. Math.*, vol. 335, Amer. Math. Soc., Providence, RI, 2003, pp. 195–207.
- [20] A. Kishimoto, A Rohlin property for one-parameter automorphism groups, *Comm. Math. Phys.* 179 (1996) 599–622.
- [21] K. Kodaka, T. Teruya, The Rohlin property for coactions of finite dimensional C^* -Hopf algebras on unital C^* -algebras, *J. Operator Theory* 74 (2) (2015) 329–369.
- [22] J. Kustermans, S. Vaes, Locally compact quantum groups, *Ann. Sci. Éc. Norm. Supér.* (4) 33 (6) (2000) 837–934, [http://dx.doi.org/10.1016/S0012-9593\(00\)01055-7](http://dx.doi.org/10.1016/S0012-9593(00)01055-7).
- [23] J. Kustermans, L. Vainerman, A. Van Daele, S.L. Woronowicz, Locally compact quantum groups, in: *Conference on Noncommutative Geometry and Quantum Groups*, Lecture Notes, in: *Contemp. Math.*, vol. 335, 2001.
- [24] A. Maes, A. Van Daele, Notes on compact quantum groups, *Nieuw Arch. Wiskd.* (4) 16 (1–2) (1998) 73–112.
- [25] N. Nawata, Finite group actions on certain stably projectionless C^* -algebras with the Rohlin property, *Trans. Amer. Math. Soc.* 368 (2016) 471–493.
- [26] R. Nest, C. Voigt, Equivariant Poincaré duality for quantum group actions, *J. Funct. Anal.* 258 (5) (2010) 1466–1503, <http://dx.doi.org/10.1016/j.jfa.2009.10.015>.
- [27] H. Osaka, N.C. Phillips, Crossed products by finite group actions with the Rokhlin property, *Math. Z.* 270 (1–2) (2012) 19–42.
- [28] P. Podleś, Symmetries of quantum spaces. Subgroups and quotient spaces of quantum $SU(2)$ and $SO(3)$ groups, *Comm. Math. Phys.* 170 (1) (1995) 1–20, <http://projecteuclid.org/getRecord?id=euclid.cmp/1104272946>.
- [29] M. Rørdam, E. Stormer, Classification of Nuclear C^* -Algebras. Entropy in Operator Algebras, *Encyclopaedia of Mathematical Sciences*, Springer, 2001.
- [30] L. Santiago, Crossed products by actions of finite groups with the Rokhlin property, *Int. J. Math.* 26 (7) (2015).
- [31] T. Timmermann, An Invitation to Quantum Groups and Duality: From Hopf Algebras to Multiplicative Unitaries and Beyond, *EMS Textbooks in Mathematics*, European Mathematical Society (EMS), Zürich, 2008.

- [32] A.S. Toms, W. Winter, Strongly self-absorbing C^* -algebras, *Trans. Amer. Math. Soc.* 359 (8) (2007) 3999–4029.
- [33] S. Vaes, R. Vergnioux, The boundary of universal discrete quantum groups, exactness, and factoriality, *Duke Math. J.* 140 (1) (2007) 35–84, <http://dx.doi.org/10.1215/S0012-7094-07-14012-2>.
- [34] S.L. Woronowicz, Compact quantum groups, in: *Symétries Quantiques*, Les Houches, 1995, North-Holland, Amsterdam, 1998, pp. 845–884.